

Radiation Beyond Four Space-Time Dimensions

A. Mironov*

Lebedev Physics Institute and ITEP, Moscow, Russia

A. Morozov†

ITEP, Moscow, Russia

Abstract

We present a list of formulas describing classical radiation of the rank s tensor field from an accelerated point-like source in flat space-time of arbitrary even dimension d . This allows straightforward evaluating the total intensity and radiated momentum for any s and d algorithmically, by hands or with the help of a computer (e.g. with an attached MAPLE program). Practical application of formulas is limited, because, for $s > 1$, the energy-momentum tensor for the point-like source is not conserved. This usually means that one cannot neglect contributions to radiation from tensions of the forces that cause acceleration of the source.

Traditionally radiation processes were considered as subjects of direct physical application and therefore were deeply investigated only in no more than $d = 4$ space-time dimensions, see [1]. Even for $d = 2$ and 3, where obvious applications exist, say, to sound waves in media, the theory remains badly represented in the literature. Only recently, after the strings-inspired multi-dimensional models [2] attracted increasing attention [3], some papers on multi-dimensional radiation began to emerge [4, 5, 6, 8, 7]. Of course, they are still too few to cover the field at the level of exhaustiveness, typical for the literature on $4d$ radiation. In this paper we make a step, which we think is necessary for a *systematic* study of physical effects. Namely, we provide a list of generic formulas, describing *classical* radiation for arbitrary dimension d and rank s of the *radiated* fields, what should help to reveal the underlying physical and mathematical structures. In particular, one can immediately read off the radiation damping force in higher dimensions from our results, for example by the method of [4].

To illustrate the main results of this paper, we use several **Tables**. We begin with **Table I**, summarizing the logic of radiation calculus. The left column is our main line, the right column enumerates related subjects which deserve deeper physical consideration and are left for a detailed presentation of radiation physics in a more sophisticated review article.

According to **Table I**, in order to find radiation intensity one needs to perform the following chain of calculations:

- Solve wave equation for a point-like source of the rank- s field, moving along a world line $z^\mu(\tau)$,

$$\square A_{\mu_1 \dots \mu_s}(x) = \oint u_{\mu_1} \dots u_{\mu_s} \delta^{(d)}(x - z(\tau)) d\tau, \quad (1)$$

and pick up the slowest-falling contribution at large distances. For even d it is given by a simple formula for the retarded Liénard-Wiechert potential:

$$A_{\mu_1 \dots \mu_s}^{rad} = \left(\frac{1}{(Ru)} \partial_\tau \right)^{\frac{d-4}{2}} \frac{u_{\mu_1} \dots u_{\mu_s}}{Ru} \quad \text{and} \quad \partial_\mu A_{\mu_1 \dots \mu_s}^{rad} = R_\mu \left(\frac{1}{(Ru)} \partial_\tau \right)^{\frac{d-2}{2}} \frac{u_{\mu_1} \dots u_{\mu_s}}{Ru} \quad (2)$$

where $u_\mu = \partial_\tau z_\mu$ is source's d -velocity, $u^2 = 1$, evaluated at the radiation-time moment $t' = z^0(\tau)$ which is determined by the condition $R^2 = 0$, where R_μ is the d -vector with components $R^\mu \equiv x^\mu - z^\mu(\tau)$. Here τ is source's proper time. In particular,

$$\frac{\partial \tau}{\partial x^\mu} = \frac{R_\mu}{(Ru)} \quad (3)$$

*E-mail: mironov@lpi.ru; mironov@itep.ru

†E-mail: morozov@itep.ru

We also introduce the notation $n^\mu \equiv (1, -\vec{n}) = \frac{R^\mu}{R}$, where R is length of the spatial part of R^μ , $R \equiv \sqrt{(R^i)^2}$ (n^μ is **not** a d -vector in contrast to R^μ ; we define it in the laboratory frame).

It is convenient to re-write (2) in condensed notation:

$$A_{rad} = \frac{1}{R^{d-2}} \left(\frac{1}{U} \partial_\tau \right)^{p-1} \frac{S}{U} = \frac{1}{R^{d-2}} \sum_{q=0}^{p-1} (\partial_\tau^q S) A_q^{p-1} \quad \text{and} \quad \partial_\mu A_{rad} = \frac{n_\mu}{R^{d-2}} \sum_{q=0}^p (\partial_\tau^q S) A_q^p, \quad (4)$$

where $U = (nu)$, $S = u_{\mu_1} \dots u_{\mu_s}$, $p = \frac{d-2}{2}$, and find A_q^p from recurrent relations

$$A_q^{p+1} = \frac{1}{U} \left(\partial_\tau A_q^p + A_{q-1}^p \right) \quad (5)$$

The first few A_q^p are listed in **Table II**.

- Check transversality: in the leading order in $1/R$, when derivatives of R are neglected,

$$\partial^{\mu_s} A_{\mu_1 \mu_2 \dots \mu_s}^{rad} = \left(\frac{1}{(Ru)} \partial_\tau \right)^{\frac{d-2}{2}} u_{\mu_1} \dots u_{\mu_{s-1}} \quad (6)$$

does not vanish for $s \geq 2$, and this is *not* cured by subtraction of traces. The physical reason for non-transversality is the disregard of radiation from tensions of the forces that cause acceleration of the source: only taking into account all the tensions makes radiation problem well-defined for $s \geq 2$. For $s \geq 2$ the formulas of the present paper provide only a part of the full answer.

- Build up the energy-momentum tensor in the wave zone. It is equal to

$$T_{\mu\nu}^{(s)} = R_\mu R_\nu \left\{ \left(\frac{1}{(Ru)} \partial_\tau \right)^{\frac{d-2}{2}} \frac{u_{\mu_1} \dots u_{\mu_s}}{Ru} \right\}^2 = \frac{R_\mu R_\nu}{R^d} \left\{ \sum_{q=0}^p A_q^p \partial_\tau^q S \right\}^2 \quad (7)$$

In applications, certain linear combinations of such stress tensors arise. For instance, for the scalar waves

$$T_{\mu\nu}^{(scalar)} = -T_{\mu\nu}^{(0)} \quad (8)$$

This is because only the spatial components of all non-zero spin fields have any physical meaning (e.g. survive in physical gauges), thus giving rise to the overall minus sign of the kinetic part of the energy-momentum tensor as compared with the scalar case. For gravitational waves

$$T_{\mu\nu}^{(grav)} = T_{\mu\nu}^{(2)} - \frac{1}{2} T_{\mu\nu}^{(0)} \quad (9)$$

Similar redefinitions are also needed for higher spins $s > 2$. These linear transformations can be easily made using the results for $T_{\mu\nu}^{(s)}$ read off from **Table V**.

- Evaluate the radiated momentum flux through the sphere of radius R . It is equal to

$$d\mathcal{P}_{d|s}^\mu = - \oint \{ T^{\mu i} n_i dx^0 \} dS \quad (10)$$

where dS is an infinitesimal element of this sphere (i.e. integration runs over the sphere). Then, one finally obtains (using $dx^0 = U d\tau = (nu) d\tau$)

$$\begin{aligned} \frac{d\mathcal{P}_{d|s}^\mu}{dt} &= - \oint T^{\mu i} n_i R^{d-2} U d\Omega_{d-2} = \\ &= \sum_{q', q''=0}^p \left(\partial_\tau^{q'} (u_{\mu_1} \dots u_{\mu_s}) \partial_\tau^{q''} (u^{\mu_1} \dots u^{\mu_s}) \right) \int n^\mu U A_{q'}^p A_{q''}^p d\Omega_{d-2} (n) = \sum_{k=0}^{\frac{d-4}{2}} (\partial_\tau^k u^\mu) P_k^{(d|s)}(\kappa) \end{aligned} \quad (11)$$

where $d\Omega_{d-2}$ is the solid angle in the $d-1$ space and $P_k^{(d|s)}$ are certain functions of various scalar products $(\partial_\tau^l u_\nu \partial_\tau^m u^\nu)$ with $l+m \leq d-4-k$. In particular, the radiated energy loss is $(dt' = u^0 d\tau = \gamma d\tau)$

$$\frac{d\mathcal{P}_{d|s}^0}{dt'} = \frac{1}{\gamma} \sum_{k=0}^{\frac{d-4}{2}} (\partial_\tau^k \gamma) P_k^{(d|s)}(\kappa) \xrightarrow{\text{for constant } \gamma=(1-v^2)^{-1/2}} P_0^{(d|s)}(\kappa), \quad (12)$$

We warn once again that due to effects like (6) and (9)-(8), this quantity represents only a part of the full answer for $s \geq 2$. Moreover, as follows from **Table V**, it can be even negative (!): for example, for fields with asymptotically large rank s the leading contribution to the radiated momentum, $\frac{d\mathcal{P}_{d|s}^\mu}{d\tau}$ behaves as $-(s\dot{u}^2)^{\frac{d}{2}-1} \sim (-)^{\frac{d}{2}}$, provided $\dot{u}^2 \leq 0$. For $s = 0$, this is cured by changing the common sign, see (8). However, for $s \geq 3$ the procedure has to be more sophisticated.

Sometime also contractions $I_{d|s} = u^\mu \frac{d\mathcal{P}_{d|s}^\mu}{d\tau}$ are considered [6], but they do not have any direct physical meaning unless $\frac{d\mathcal{P}_{d|s}^\mu}{d\tau} \sim u_\mu$, as happens for $d = 4$. In this latter case, $I_{d|s} = \frac{d\mathcal{P}_{d|s}^0}{dt}$.

- Calculate angular integrals over isotropic (light-like) unit vectors n^μ in (11) (although n^μ is not a d -vector, these integrals are Lorentz-covariant, since (11) is Lorentz-covariant)),

$$\int \frac{n^{\mu_1} \dots n^{\mu_m}}{(nu)^{d+m-2}} d\Omega_{d-2}(n) \sim \text{Pr}_d(u^{\mu_1} \dots u^{\mu_m}) \quad (13)$$

They are expressed through the trace-eliminating projector Pr_d (since $n^2 = 0$ and contraction of any pair of μ -indices at the l.h.s. should give zero):

$$\begin{aligned} \text{spin } s = 2 : & \quad \text{Pr}_d(u_\mu u_\nu) = u_\mu u_\nu - \frac{1}{d} \eta_{\mu\nu} \\ \text{spin } s = 3 : & \quad \text{Pr}_d(u_{\mu_1} u_{\mu_2} u_{\mu_3}) = u_{\mu_1} u_{\mu_2} u_{\mu_3} - \frac{1}{d+2} \left(\eta_{\mu_1 \mu_2} u_{\mu_3} + \eta_{\mu_1 \mu_3} u_{\mu_2} + \eta_{\mu_2 \mu_3} u_{\mu_1} \right) \\ \text{spin } s = 4 : & \quad \text{Pr}_d(u_{\mu_1} u_{\mu_2} u_{\mu_3} u_{\mu_4}) = u_{\mu_1} u_{\mu_2} u_{\mu_3} u_{\mu_4} - \\ & \quad - \frac{1}{d+4} \left(\eta_{\mu_1 \mu_2} u_{\mu_3} u_{\mu_4} + 5 \text{ permutations} \right) + \frac{1}{(d+4)(d+2)} \left(\eta_{\mu_1 \mu_2} \eta_{\mu_3 \mu_4} + \eta_{\mu_1 \mu_3} \eta_{\mu_2 \mu_4} + \eta_{\mu_1 \mu_4} \eta_{\mu_2 \mu_3} \right) \\ & \quad \dots \end{aligned}$$

For generic m we have:

$$\text{Pr}_d(u_{\mu_1} \dots u_{\mu_m}) = \sum_{k=0}^{[m/2]} (-)^k \frac{(d+2m-4-2k)!!}{(d+2m-4)!!} \left(\overbrace{u_{\mu_1} \dots u_{\mu_{m-2k}} \left(\underbrace{\eta_{\mu_{m-2k+1} \mu_{m-2k+2}} \dots \eta_{\mu_{m-1} \mu_m} + \dots}_{(2k-1)!! \text{ permutations}} \right)}^{C_{2k}^m \text{ permutations}} + \dots \right)$$

This projector can be conveniently re-written through generating function: after contraction with m copies of a d -vector x we get

$$\begin{aligned} \text{Pr}_d((xu)^m) &= \sum_{k=0}^{[m/2]} (-)^k (2k-1)!! C_{2k}^m \frac{(d+2m-4-2k)!!}{(d+2m-4)!!} (xu)^{m-2k} (x^2)^k = \\ &= \sum_{k=0}^{[m/2]} (-)^k \frac{(d+2m-4-2k)!!}{(d+2m-4)!!} \frac{m! (xu)^{m-2k} (x^2)^k}{2^k k! (m-2k)!} \end{aligned} \quad (14)$$

The sum of coefficients in this formula is needed to define the normalization factor in (13):

$$c_d(m) = \sum_{k=0}^{[m/2]} (-)^k C_{2k}^m (2k-1)!! \frac{(d+2m-4-2k)!!}{(d+2m-4)!!} = \frac{(d+2[m/2]-3)!! (d+2(m-[m/2])-4)!!}{(d-3)!! (d+2m-4)!!} \quad (15)$$

Here $[m/2]$ means integer part of a number, e.g. $[3/2] = 1$ and $3 - [3/2] = 2$.

Rewritten in terms of the generating function and with normalization factor restored, the angular integral, (13) is (see **Table III**):

$$\begin{aligned} \int \frac{(nx)^m d\Omega_{d-2}(n)}{(nu)^{d+m-2}} &= \frac{S_{n-2}}{c_d(s)} \text{Pr}_d((ux)^m) = \\ &= \frac{2^{d/2} \pi^{\frac{d-2}{2}}}{(d+2[m/2]-3)!!} \sum_{j=0}^{[m/2]} (-)^j \frac{(d+2m-4-2j)!!}{(d+2m-2[m/2]-4)!!} \frac{m!}{2^j j! (m-2j)!} (ux)^{m-2j} (x^2)^j \end{aligned} \quad (16)$$

It remains to substitute the vector x by $x = tx_0 + \sum_{k=0}^{\frac{d-2}{2}} a_k \partial_\tau^k u$ and pick up the coefficient in front of the relevant combination of t and a_k . Note that this calculation uses the generation function which provides an

additional combinatorial coefficient that has to be taken into account manifestly. E.g., in order to get the correct coefficients in front of the term $\partial_\tau^{k_1} u \dots \partial_\tau^{k_n} u$ with different values of $k_1, \dots, k_n$, one has to divide the generating function by $n!$.

The final results can be equally expressed via scalar products $U_{ij} \equiv \partial_\tau^l \vec{u}_\nu \partial_\tau^m \vec{u}^\nu$ and through the Frenet curvatures κ_m and their derivatives $\partial_\tau^l \kappa_m$. Hence, the last step:

- Re-express U_{ij} through Frenet curvatures [10] which parameterize the moving orthonormal basis, associated with the world line $x^\mu(\tau)$. It is formed by d -vectors $\vec{N}^{(\mu)}$, $\mu = 0, \dots, d-1$,

$$\vec{N}^{(\mu)} \vec{N}^{(\nu)} = \eta^{\mu\nu} \quad (17)$$

with $\vec{N}^{(0)} = \vec{u}$ (i.e. $N_\mu^{(0)} = u_\mu$) and other vectors constructed recursively from the condition

$$\partial_\tau \vec{N}^{(\mu)} = -k_{\mu+1} \vec{N}^{(\mu+1)} + \sum_{\nu=1}^{\mu} \beta_{\mu\nu} \vec{N}^{(\nu)}$$

From τ -derivative of orthonormality condition,

$$\left(\partial_\tau \vec{N}^{(\mu)} \right) \vec{N}^{(\nu)} + \vec{N}^{(\mu)} \left(\partial_\tau \vec{N}^{(\nu)} \right) = 0 .$$

It follows that at $\nu \leq \mu$

$$\beta_{\mu\nu} = \left(\partial_\tau \vec{N}^{(\mu)} \right) \vec{N}^{(\nu)} = -\vec{N}^{(\mu)} \left(\partial_\tau \vec{N}^{(\nu)} \right) = \kappa_\mu \delta_{\nu, \mu-1} \eta^{\mu\mu}$$

and

$$\partial_\tau \vec{N}^{(\mu)} = -k_{\mu+1} \vec{N}^{(\mu+1)} + k_\mu \vec{N}^{\mu-1} \quad (18)$$

Parameters k_μ (they are not d -vectors!) depend on the shape of the world line $x^\mu(\tau)$ in the infinitesimal vicinity of its point and are called Frenet curvatures. Expressions of scalar products U_{ij} through these curvatures are given in **Table IV**.

Results

All these steps are easily made with the help of MAPLE or Mathematica, see Appendix at the end of this paper for a MAPLE program, [9]. As an application of the algorithm we obtain the explicit formulas for the radiated d -momentum $d\mathcal{P}_{d|s}^\mu$ as well as somewhat simpler formulas for $P_0^{(d|s)}$, the radiated intensity at constant γ , (i.e.

$\kappa_2^2 = \frac{\gamma^2 \kappa_1^2}{\gamma^2 - 1} = \text{const}$, $\kappa_i = 0$, $i > 2$) see (12), and for the contractions $I_{d|s} = u^\mu \frac{d\mathcal{P}_{d|s}^\mu}{d\tau}$, [6] for “realistic” values of $d = 4, 6, 8, 10$ and generic rank s . First, we list results for the case of $d = 4, 6, 8$, when the formulas are relatively simple, and one can understand their general structure. Much more involved formulas in the most interesting case of $d = 10$ are put into a separate section.

$$\boxed{d\mathcal{P}_{d|s}^\mu}$$

$$\frac{d\mathcal{P}_{4|s}^\mu}{d\tau} = \frac{4-12s}{3}\pi\dot{u}^2u^\mu$$

$$\frac{d\mathcal{P}_{6|s}^\mu}{d\tau} = \pi^2 \left[\frac{19}{3} - (2s-3)^2 \right] \dot{u}^4 u^\mu + \frac{8\pi^2}{15}(1-5s)\ddot{u}^2 u^\mu + \frac{16\pi^2}{35}(2-7s)(\dot{u}\ddot{u})\dot{u}^\mu + \frac{16\pi^2}{105}(7s-4)\dot{u}^2\ddot{u}^\mu$$

$$\begin{aligned} \frac{d\mathcal{P}_{8|s}^\mu}{d\tau} = & \left\{ -\frac{16\pi^3}{9} \left(\frac{7s^3}{10} - \frac{135}{14}s^2 + \frac{838}{35}s - 5 \right) \dot{u}^6 + \frac{32\pi^3}{3} \left(-s^2 + \frac{101}{35}s - \frac{41}{63} \right) (\dot{u}\ddot{u})^2 + \right. \\ & + \frac{16\pi^3}{3} \left(-\frac{s^2}{5} + \frac{59}{35}s - \frac{20}{63} \right) \dot{u}^2\ddot{u}^2 + 32\pi^3 \left(\frac{s^2}{15} - \frac{2s}{7} + \frac{11}{189} \right) \dot{u}^2(\dot{u}\ddot{u}) - \frac{16\pi^3}{15} \left(s - \frac{1}{7} \right) \dot{\ddot{u}}^2 \Big\} u^\mu + \\ & + \left\{ \frac{32\pi^3}{7} \left(-\frac{6s^2}{5} + \frac{19}{5}s - \frac{283}{297} \right) \dot{u}^2(\dot{u}\ddot{u}) + \frac{64\pi^3}{7} \left(\frac{1}{27} - \frac{s}{5} \right) \ddot{u}\dot{\ddot{u}} \right\} \dot{u}^\mu + \\ & + \left\{ \frac{64\pi^3}{21} \left(\frac{2s^2}{5} - 2s + \frac{65}{99} \right) \dot{u}^4 + \frac{64\pi^3}{21} \left(\frac{1}{9} - \frac{2s}{5} \right) \dot{u}\dot{\ddot{u}} \right\} \ddot{u}^\mu + \frac{32\pi^3}{35} \left(s - \frac{17}{27} \right) (\dot{u}\ddot{u})\dot{\ddot{u}}^\mu \end{aligned}$$

$$\boxed{P_0^{(d|s)}}$$

$$P_0^{(4|s)} = \frac{12s-4}{3}\pi\kappa_1^2$$

$$P_0^{(6|s)} = -\pi^2 \left(4s^2 - \frac{28}{3}s + \frac{32}{15} \right) 4\kappa_1^4 + \frac{8\pi^2}{15}(5s-1)\kappa_1^2\kappa_2^2$$

$$P_0^{(8|s)} = \frac{16\pi^3}{105}(7s-1)(\kappa_1^2\kappa_2^2\kappa_3^2 + \kappa_1^2\kappa_2^4) + \frac{8\pi^3}{315}(49s^3 - 549s^2 + 1004s - 216)\kappa_1^6 - 16\pi^3 \left(\frac{s^2}{5} - s + \frac{64}{315} \right) \kappa_1^4\kappa_2^2$$

$$\boxed{I_{d|s}}$$

$$I_{4|s} = \frac{12s-4}{3}\pi\kappa_1^2$$

$$I_{6|s} = -\pi^2 \left(4s^2 - \frac{124}{15}s + \frac{32}{21} \right) \kappa_1^4 + \frac{8\pi^2}{15}(5s-1)(\dot{\kappa}_1^2 + \kappa_1^2\kappa_2^2)$$

$$\begin{aligned} I_{8|s} = & \frac{16\pi^3}{105} \left[\left(\frac{49}{6}s^3 - \frac{167}{2}s^2 + \frac{406}{3}s - \frac{276}{11} \right) \kappa_1^6 - \left(21s^2 - 97s + \frac{172}{9} \right) \kappa_1^4\kappa_2^2 - \left(77s^2 - 180s + \frac{109}{3} \right) \kappa_1^2\dot{\kappa}_1^2 + \right. \\ & \left. + 2(7s^2 - 9s + 4)\kappa_1^3\ddot{\kappa}_1 + (7s-1) \left\{ \kappa_1^2\kappa_2^2\kappa_3^2 + (\kappa_1\kappa_2^2 - \ddot{\kappa}_1)^2 + (2\kappa_2\dot{\kappa}_1 + \kappa_1\dot{\kappa}_2)^2 \right\} \right] \end{aligned}$$

Results for the most interesting case of $d = 10$

$$\begin{aligned}
\frac{dP_{10|s}^\mu}{d\tau} = & \left\{ \frac{4\pi^4}{315} \left(19s^4 - \frac{6466}{9}s^3 + \frac{236237}{33}s^2 - \frac{1609666}{99}s + \frac{9800}{3} \right) \dot{u}^8 - \frac{32\pi^4}{7} \left(s^3 - \frac{200}{9}s^2 + \frac{2147}{33}s - \frac{6230}{429} \right) \dot{u}^2(\dot{u}\ddot{u})^2 + \right. \\
& + \frac{16\pi^4}{35} \left(s^3 - \frac{61}{27}s^2 - \frac{8316}{297}s + \frac{11690}{1287} \right) \dot{u}^4\ddot{u}^2 + \frac{64\pi^4}{35} \left(s^3 - \frac{1037}{54}s^2 + \frac{32653}{594}s - \frac{4865}{429} \right) \dot{u}^4(\dot{u}\ddot{u}) - \\
& - \frac{16\pi^4}{105} \left(19s^2 - \frac{611}{9}s + \frac{512}{33} \right) \ddot{u}^4 - \frac{32\pi^4}{15} \left(\frac{17}{7}s^2 - 151s + \frac{34}{21} \right) (\dot{u}\ddot{u})^2 - \frac{64\pi^4}{105} \left(s^2 - \frac{53}{3}s + \frac{100}{33} \right) (\dot{u}\ddot{u})(\ddot{u}\ddot{u}) - \\
& - \frac{256\pi^4}{35} \left(s^2 - \frac{263}{108}s + \frac{119}{198} \right) \ddot{u}^2(\dot{u}\ddot{u}) - \frac{32\pi^4}{105} \left(s^2 - \frac{134}{9}s + \frac{25}{11} \right) \dot{u}^2\ddot{u}^2 + \frac{64\pi^4}{105} \left(s^2 - \frac{74}{9}s + \frac{15}{11} \right) \dot{u}^2(\ddot{u}\ddot{u}) + \\
& + \frac{64\pi^4}{315} (9s^2 - 49s + 10)(\dot{u}\ddot{u})(\ddot{u}\ddot{u}) + \frac{32\pi^4}{105} \left(\frac{1}{9} - s \right) \ddot{u}^4 \Big\} u^\mu - \left\{ \frac{32\pi^4}{63} \left(\frac{8s^3}{3} - \frac{593}{11}s^2 + \frac{72173}{429}s - \frac{5670}{143} \right) \dot{u}^4(\dot{u}\ddot{u}) + \right. \\
& + \frac{64\pi^4}{189} \left(20s^2 - \frac{829}{11}s + \frac{2664}{143} \right) \ddot{u}^2(\dot{u}\ddot{u}) + \frac{256\pi^4}{189} \left(5s^2 - \frac{133}{11}s + \frac{453}{143} \right) (\dot{u}\ddot{u})(\dot{u}\ddot{u}) - \frac{128\pi^4}{189} \left(s^2 - \frac{109}{22}s + \frac{147}{143} \right) \dot{u}^2(\ddot{u}\ddot{u}) + \\
& + \frac{320\pi^4}{819} \left(1 - \frac{208}{33}s \right) \dot{u}^2(\ddot{u}\ddot{u}) + \frac{64\pi^4}{189} \left(2s - \frac{3}{11} \right) (\ddot{u}\ddot{u}) \Big\} \dot{u}^\mu + \left\{ \frac{128\pi^4}{189} \left(s^3 - \frac{785}{44}s^2 + \frac{30843}{572}s - \frac{2065}{143} \right) \dot{u}^6 + \right. \\
& + \frac{320\pi^4}{189} \left(s^2 - \frac{53s}{11} + \frac{216}{143} \right) (\dot{u}\ddot{u})^2 - \frac{640\pi^4}{189} \left(s^2 - 4s + \frac{1569}{1430} \right) \dot{u}^2(\dot{u}\ddot{u}) - \frac{64\pi^4}{189} \left(4s^2 + s + \frac{123}{143} \right) \dot{u}^2\ddot{u}^2 + \\
& + \frac{128\pi^4}{189} \left(\frac{2}{11} - s \right) (\ddot{u}\ddot{u}) \Big\} \ddot{u}^\mu + \left\{ \frac{256\pi^4}{4158} \left(22s^2 - 167s + \frac{795}{13} \right) \dot{u}^2(\dot{u}\ddot{u}) + \frac{64\pi^4}{189} \left(\frac{3}{11} - s \right) (\dot{u}\ddot{u}) \right\} \ddot{u}^\mu - \\
& - \left\{ \frac{32\pi^4}{945} \left(s^2 - \frac{141}{11}s + \frac{1020}{143} \right) \dot{u}^4 - \frac{64\pi^4}{315} \left(s - \frac{23}{33} \right) \ddot{u}^2 - \frac{256\pi^4}{945} \left(s - \frac{29}{44} \right) (\dot{u}\ddot{u}) \right\} \ddot{u}^\mu
\end{aligned}$$

$$\begin{aligned}
P_0^{(10|s)} = & \frac{32\pi^4}{315} \left[\left(-\frac{19}{8}s^4 + \frac{2747}{36}s^3 - \frac{152021}{264}s^2 + \frac{369827}{396} - 192 \right) \kappa_1^8 + \left(\frac{9}{2}s^3 - 605s^2 + 933s - 188 \right) \kappa_1^6\kappa_2^2 - \right. \\
& - \left(\frac{33}{2}s^2 - 325s + 28 \right) \kappa_1^4\kappa_2^4 - \left(9s^2 - \frac{88}{3}s + \frac{43}{3} \right) \kappa_1^4\kappa_2^2\kappa_3^2 + \left(3s - \frac{1}{3} \right) (\kappa_1^2\kappa_2^6 + \kappa_1^2\kappa_2^2\kappa_3^2\kappa_4^2 + 2\kappa_1^2\kappa_2^4\kappa_3^2 + \kappa_1^2\kappa_2^2\kappa_3^4) \Big]
\end{aligned}$$

$$I_{10|s} = \frac{16\pi^4}{945} \left[\left(81s^3 - 1693s^2 + \frac{53244}{11}s - \frac{133048}{143} \right) \kappa_1^6\kappa_2^2 + \left(243s^3 - 4785s^2 + \frac{106818}{11}s - \frac{300280}{143} \right) \kappa_1^4\dot{\kappa}_1^2 - \right.$$

$$\begin{aligned}
& - \left(99s^2 - 931s + \frac{1744}{11} \right) \kappa_1^4 \kappa_2^4 + \left(36s^2 - 220s + \frac{416}{11} \right) \kappa_1^4 \kappa_2 \ddot{\kappa}_2 + \left(324s^2 - 1780s + \frac{3392}{11} \right) \kappa_1^3 \ddot{\kappa}_1 \kappa_2^2 - \\
& - \left(468s^2 - 1160s + \frac{2812}{11} \right) \kappa_1 \ddot{\kappa}_1 \kappa_1^2 + \left(144s^2 - 568s + \frac{1336}{11} \right) \kappa_1^2 \dot{\kappa}_1 \dot{\kappa}_1 - \left(171s^2 - 413s + \frac{1062}{11} \right) \dot{\kappa}_1^4 - \\
& - \left(54s^2 - 488s + \frac{866}{11} \right) \kappa_1^4 \kappa_2^2 \kappa_3^2 - \left(\frac{57}{4}s^4 - \frac{2507}{6}s^3 + \frac{125989}{44}s^2 - \frac{3644975}{858}s + \frac{111744}{143} \right) \kappa_1^8 - \\
& - \left(108s^3 - 1506s^2 + \frac{31794}{11}s - \frac{83808}{143} \right) \kappa_1^5 \ddot{\kappa}_1 - \left(324s^2 - 822s + \frac{1878}{11} \right) \kappa_1^2 \ddot{\kappa}_1^2 - \left(432s^2 - 2752s + \frac{5760}{11} \right) \kappa_1^3 \dot{\kappa}_1 \kappa_2 \dot{\kappa}_2 - \\
& - \left(18s^2 - 268s + \frac{450}{11} \right) \kappa_1^4 \dot{\kappa}_2^2 - \left(450s^2 - 3714s + \frac{7520}{11} \right) \kappa_1^2 \dot{\kappa}_1^2 \kappa_2^2 + \\
& + 2(9s - 1)[(\dot{\kappa}_1 - 3\dot{\kappa}_1 \kappa_2^2 - 3\kappa_1 \kappa_2 \dot{\kappa}_2)^2 + (3\ddot{\kappa}_1 \kappa_2 + 3\dot{\kappa}_1 \dot{\kappa}_2 + \kappa_1 \ddot{\kappa}_2 - \kappa_1 \kappa_2^3)^2 + (2\kappa_1 \dot{\kappa}_2 \kappa_3 + \kappa_1 \kappa_2 \dot{\kappa}_3 + 3\dot{\kappa}_1 \kappa_2 \kappa_3)^2 - \\
& - \kappa_1^2 \kappa_2^2 \kappa_3^2 (2\kappa_2^2 + \kappa_3^2 + \kappa_4^2) - 2\kappa_1 \kappa_2 \kappa_3^2 (3\dot{\kappa}_1 \kappa_2 + \kappa_1 \dot{\kappa}_2 + 3\dot{\kappa}_1 \dot{\kappa}_2)]
\end{aligned}$$

Note that, in the expressions for the radiated momentum the coefficients in front of terms of degree k in U_{ij} are polynomials of degree k in rank s . Relative simplicity of the coefficients of these polynomials implies that a general formula may exist for them at any dimension d and for arbitrary rank s . However, we stress once again that these expressions for higher ranks are of restricted use, because of necessity to take into account contributions from other energy-momentum tensors. The second thing one has to do in order to separate the contribution of the given *spin* s from the expression for the *rank* s is to add proper combinations of energy-momentum tensors for lower rank fields. This is done straightforward using formulas we obtained here.

The examples of these formulas, with illustrative purposes, are collected in **Table V** for low values of $s = 0, 1, 2, 3$ for dimensions $d = 4, 6, 8$ when the formulas are more comprehensible. Much more involved (but interesting) case of $d = 10$ is listed after that table. The Appendix contains an example of the MAPLE program for calculating various quantities obtained in this paper.

Acknowledgements

We appreciate informative discussions with P.Kazinsky and S.Lyakhovich. Our work is partly supported by the program of the Federal Nuclear Energy Agency, by grants RFBR 07-02-00878- (A.Mir.) and 07-02-00645 (A.Mor.), joint grant 06-01-92059-CE, by the Grant of Support for the Scientific Schools NSH-8004.2006.2, by NWO project 047.011.2004.026, INTAS grant 05-1000008-7865 and ANR-05-BLAN-0029-01 project.

References

- [1] L.Landau and E.Lifshitz, *The Classical Theory of Fields (Course of Theoretical Physics, Volume 2)* J.D.Jackson, *Classical Electrodynamics*, Wiley, New York, 1975
A.A.Sokolov, I.M.Ternov, V.Ch.Zhukovsky and A.B.Borisov, *Quantum Electrodynamics*, Moscow State University Press, 1983
S.Weinberg, *Gravitation and Cosmology*, New York: Wiley, 1972
I.Khriplovich, *General Relativity* (in Russian), Izhevsk, 2001

- [2] N.Arkani-Khamed, S.Dimopoulos and G.Dvali, Phys.Lett. **B429** (1998) 263; Phys.Rev. **D59** (1999) 086004
I.Antoniadis, N.Arkani-Khamed, S.Dimopoulos and G.Dvali, Phys.Lett. **B436** (1998) 257
- [3] T.Banks and W.Fishler, hep-th/9906038
S.B.Giddings and S.Thomas, Phys.Rev. **D65** (2002) 056010
S.Dimopoulos and G.Landsberg, Phys.Rev.Lett. **87** (2001) 161602
References and a review can be found in:
P.Kanti, Int.J.Mod.Phys. **A19** (2004) 4899
G.Landsberg, J.Phys. **G32** (2006) R337, hep-ph/0607297
M.Cavaglia, R.Godang, L.Cremaldi and D.Summers, hep-ph/0609001
For mini-black-holes in cosmic ray events, see:
A.Mironov, A.Morozov and T.Tomaras, Sov.J.Nucl.Phys. (Yad.Fiz.), hep-ph/0311318
A.Cafarella, C.Coriano and T.Tomaras, JHEP **0506** (2005) 065, hep-th/0410358
For mini-black-holes in neutrino experiments, see a very detailed review in:
L.Anchordoqui, T.Paul, S.Reucroft and J.Swain, Int.J.Mod.Phys. **A18** (2003) 2229
and references therein
For more exotic possibilities, see:
A.Mironov, A.Morozov and T.Tomaras, If LHC is a Mini-Time-Machines Factory, Can We Notice? Proceedings of the II Southeastern European Workshop "Challenges Beyond the Standard Model", 2005
- [4] B.Kosyakov, Theor.Math.Phys. **119** (1999) 493-505, hep-th/0207217
- [5] D.Galtsov, Phys.Rev. **D66** (2002) 025016, hep-th/0112110
P.Kazinsky, S.Lyakhovich and A.Sharapov, Phys.Rev. **D66** (2002) 025017, hep-th/0201046
V.Cardoso, O.Dias and J.Lemos, Phys.Rev. **D67** (2003) 064026, hep-th/0212168
Yu.Yaremenko, J.Phys. **A37** (2004) 1079-1091
B.Koch and M.Bleicher, hep-th/0512353
P.Krtous and J.Podolsky, Class.Quant.Grav. **23** (2006) 1603-1616, gr-qc/0602007
H.A.Morales-Técotl, O.Pedraza and L.O.Pimentel, physics/0611241
- [6] M.Gurses and O.Sarioglu, *Class.Quant.Grav.* **19** (2002) 4249; **20** (2003) 351; hep-th/0303078
- [7] V.Cardoso, M.Cavaglia and J.-Q.Guo, hep-th/0702138
- [8] A.Mironov and A.Morozov, Pisma Zh.Eksp.Teor.Fiz. **85** (2007) 9-14 (JETP letters, **85** (2007) 6-11), hep-ph/0612074
- [9] <http://thesaurus.itep.ru/project/0703mm>
- [10] M.Spivak, *A Comprehensive Introduction to Differential Geometry*, Boston, 1979

Tables

TABLE I		Radiation: calculus and problems	
accelerating point-like body (source)			
↓		↘	
radiation field		separation of radiation from co-moving field, wave zone	
↓		↘	
even dimensions		strong violation of Huygens principle at finite distances in odd dimensions	
↓		↘	
retarded field	$A_{rad}^{\mu_1 \dots \mu_s} = \left(\frac{1}{(Ru)} \partial_\tau \right)^{\frac{d-4}{2}} \frac{u^{\mu_1} \dots u^{\mu_s}}{(Ru)}$	difference between observation and radiation times	
↓		↘	
field tension		transversality violation for $s > 1$	
$F_\mu^{\mu_1 \dots \mu_s} = A_\mu^{\mu_1 \dots \mu_s} \sim R_\mu \left(\frac{1}{(Ru)} \partial_\tau \right)^{\frac{d-2}{2}} \frac{u^{\mu_1} \dots u^{\mu_s}}{(Ru)}$		$F_{\mu_s}^{\mu_1 \dots \mu_s} \sim \left(\frac{1}{(Ru)} \partial_\tau \right)^{\frac{d-2}{2}} u^{\mu_1} \dots u^{\mu_{s-1}}$	other sources of radiation can not be neglected
↓		↘	
stress tensor		picking the definite-spin contribution,	
$T_{\mu\nu} = R_\mu R_\nu \left\{ \left(\frac{1}{(Ru)} \partial_\tau \right)^{\frac{d-2}{2}} \frac{u^{\mu_1} \dots u^{\mu_s}}{(Ru)} \right\}^2$		radiation from other sources to cure $T_{\mu\nu}$ non-consevation	
↓		↘	
full intensity		radiation friction	
↓			
polarization, angular and frequency distributions			

TABLE II : A_q^p	$q = 0$	$q = 1$	$q = 2$	$q = 3$	$q = 4$
field in $d = 4$ $p = 0$	$\frac{1}{U}$	—	—	—	—
field in $d = 6$ $p = 1$ tension in $d = 4$	$-\frac{\dot{U}}{U^3}$	$\frac{1}{U^2}$	—	—	—
field in $d = 8$ $p = 2$ tension in $d = 6$	$\frac{-U\ddot{U}+3\dot{U}^2}{U^5}$	$-\frac{3\dot{U}}{U^4}$	$+\frac{1}{U^3}$	—	—
field in $d = 10$ $p = 3$ tension in $d = 8$	$\frac{-\dot{U}U^2+10\ddot{U}\dot{U}U-15\dot{U}^3}{U^7}$	$\frac{-4\ddot{U}U+15\dot{U}^2}{U^6}$	$-\frac{6\dot{U}}{U^5}$	$\frac{1}{U^4}$	—
field in $d = 12$ $p = 4$ tension in $d = 10$	$\frac{-\ddot{U}U^3+15\dot{U}\dot{U}U^2+10\ddot{U}^2U^2-105\ddot{U}\dot{U}^2U+105\dot{U}^4}{U^9}$	$\frac{-5\dot{U}U^2+60\ddot{U}\dot{U}U-105\dot{U}^3}{U^8}$	$\frac{-10\ddot{U}U+45\dot{U}^2}{U^7}$	$-\frac{10\dot{U}}{U^6}$	$\frac{1}{U^5}$

	TABLE III :	Formulas (16)	for angular integrals	
m	$d = 4$ ($p = 0$)	$d = 6$ ($p = 1$)	$d = 8$ ($p = 2$)	$d = 10$ ($p = 3$)
$m = 1$	$4\pi(ux)$	$\frac{8\pi^2}{3}(ux)$	$\frac{16\pi^3}{15}(ux)$	$\frac{32\pi^4}{105}(ux)$
$m = 2$	$\frac{4\pi}{3}\left(4(ux)^2 - x^2\right)$	$\frac{8\pi^2}{15}\left(6(ux)^2 - x^2\right)$	$\frac{16\pi^3}{105}\left(8(ux)^2 - x^2\right)$	$\frac{32\pi^4}{945}\left(10(ux)^2 - x^2\right)$
$m = 3$	$\frac{4\pi}{3}\left(6(ux)^3 - 3(ux)x^2\right)$	$\frac{8\pi^2}{15}\left(8(ux)^3 - 3(ux)x^2\right)$	$\frac{16\pi^3}{105}\left(10(ux)^3 - 3(ux)x^2\right)$	$\frac{32\pi^4}{945}\left(12(ux)^3 - 3(ux)x^2\right)$
$m = 4$		$\frac{8\pi^2}{105}\left(10 \cdot 8(ux)^4 - 8 \cdot 6(ux)^2 x^2 + 3(x^2)^2\right)$	$\frac{16\pi^3}{945}\left(12 \cdot 10(ux)^4 - 10 \cdot 6(ux)^2 x^2 + 3(x^2)^2\right)$	$\frac{32\pi^4}{10395}\left(14 \cdot 12(ux)^4 - 12 \cdot 6(ux)^2 x^2 + 3(x^2)^2\right)$
$m = 5$		$\frac{8\pi^2}{105}\left(12 \cdot 10(ux)^5 - 10 \cdot 10(ux)^3 x^2 + 5 \cdot 3(ux)(x^2)^2\right)$	$\frac{16\pi^3}{945}\left(14 \cdot 12(ux)^5 - 12 \cdot 10(ux)^3 x^2 + 5 \cdot 3(ux)(x^2)^2\right)$	$\frac{32\pi^4}{10395}\left(16 \cdot 14(ux)^5 - 14 \cdot 10(ux)^3 x^2 + 5 \cdot 3(ux)(x^2)^2\right)$
$m = 6$			$\frac{16\pi^3}{10395}\left(16 \cdot 14 \cdot 12(ux)^6 - 14 \cdot 12 \cdot 15(ux)^4 x^2 + 12 \cdot 15 \cdot 3(ux)^2 (x^2)^2 - 15(x^2)^3\right)$	$\frac{32\pi^4}{135135}\left(18 \cdot 16 \cdot 14(ux)^6 - 16 \cdot 14 \cdot 15(ux)^4 x^2 + 14 \cdot 15 \cdot 3(ux)^2 (x^2)^2 - 15(x^2)^3\right)$
$m = 7$			$\frac{16\pi^3}{10395}\left(18 \cdot 16 \cdot 14(ux)^7 - 16 \cdot 14 \cdot 21(ux)^5 x^2 + 14 \cdot 35 \cdot 3(ux)^3 (x^2)^2 - 15(ux)(x^2)^3\right)$	$\frac{32\pi^4}{135135}\left(20 \cdot 18 \cdot 16(ux)^7 - 18 \cdot 16 \cdot 21(ux)^5 x^2 + 16 \cdot 35 \cdot 3(ux)^3 (x^2)^2 - 15(ux)(x^2)^3\right)$
$m = 8$				$\frac{32\pi^4}{2027025}\left(22 \cdot 20 \cdot 18 \cdot 16(ux)^8 - 20 \cdot 18 \cdot 16 \cdot 28(ux)^6 x^2 + 18 \cdot 16 \cdot 70 \cdot 3(ux)^4 (x^2)^2 - 16 \cdot 28 \cdot 15(ux)^2 (x^2)^3 + 105(x^2)^4\right)$
$m = 9$				$\frac{32\pi^4}{2027025}\left(24 \cdot 22 \cdot 20 \cdot 18(ux)^9 - 22 \cdot 20 \cdot 18 \cdot 36(ux)^7 x^2 + 20 \cdot 18 \cdot 126 \cdot 3(ux)^5 (x^2)^2 - 18 \cdot 84 \cdot 15(ux)^3 (x^2)^3 + 9 \cdot 105(ux)(x^2)^4\right)$

TABLE IV :	Scalar products through Frenet curvatures	Unit vectors $\vec{N}^{(\nu)}$, $\vec{N}^{(\mu)}\vec{N}^{(\nu)} = \eta^{\mu\nu}$
order 0	$u^2 = 1$	$\vec{N}^{(0)} = \vec{u}, \quad \dot{\vec{N}}^{(0)} = \kappa_1 \vec{N}^{(1)}$
order 1	$u\dot{u} = 0$ $\dot{u}^2 = -\kappa_1^2$	$\vec{N}^{(1)} = \frac{\dot{\vec{u}}}{\kappa_1}, \quad \dot{\vec{N}}^{(1)} = \kappa_1 \vec{N}^{(0)} - \kappa_2 \vec{N}^{(2)}$
order 2	$u\ddot{u} = \kappa_1^2$ $\dot{u}\ddot{u} = -\kappa_1 \dot{\kappa}_1$ $\ddot{u}^2 = \kappa_1^4 - \kappa_1^2 \kappa_2^2 - \dot{\kappa}_1^2$	$\vec{N}^{(2)} = \frac{1}{\kappa_1 \kappa_2} \left(-\ddot{\vec{u}} + \frac{\dot{\kappa}_1}{\kappa_1} \dot{\vec{u}} + \kappa_1^2 \vec{u} \right)$ $\dot{\vec{N}}^{(2)} = \kappa_2 \vec{N}^{(1)} - \kappa_3 \vec{N}^{(3)}$
order 3	$u\dot{\dot{u}} = 3\kappa_1 \dot{\kappa}_1$ $\dot{u}\dot{\dot{u}} = -\kappa_1 \ddot{\kappa}_1 - \kappa_1^4 + \kappa_1^2 \kappa_2^2$ $\ddot{u}\dot{\dot{u}} = -\dot{\kappa}_1 \ddot{\kappa}_1 + 2\kappa_1^3 \dot{\kappa}_1 - \kappa_1 \kappa_2^2 \dot{\kappa}_1 - \kappa_1^2 \kappa_2 \dot{\kappa}_2$ $\dot{\dot{u}}^2 = -\ddot{\kappa}_1^2 - 2\kappa_1^3 \ddot{\kappa}_1 + 2\kappa_1 \kappa_2^2 \ddot{\kappa}_1 + 9\kappa_1^2 \dot{\kappa}_1^2 - 4\kappa_2^2 \dot{\kappa}_1^2 - 4\kappa_1 \kappa_2 \dot{\kappa}_1 \dot{\kappa}_2 - \kappa_1^2 \dot{\kappa}_2^2 - \kappa_1^6 + 2\kappa_1^4 \kappa_2^2 - \kappa_1^2 \kappa_2^4 - \kappa_1^2 \kappa_2^2 \kappa_3^2$	$\vec{N}^{(3)} = \frac{1}{\kappa_1 \kappa_2 \kappa_3} \left(\ddot{\vec{u}} - \left(\frac{2\dot{\kappa}_1}{\kappa_1} + \frac{\dot{\kappa}_2}{\kappa_2} \right) \dot{\vec{u}} + \left(\kappa_2^2 - \kappa_1^2 + \frac{2\dot{\kappa}_1^2}{\kappa_1^2} + \frac{\dot{\kappa}_1 \dot{\kappa}_2}{\kappa_1 \kappa_2} - \frac{\ddot{\kappa}_1}{\kappa_1} \right) \vec{u} + \kappa_1^2 \left(\frac{\dot{\kappa}_2}{\kappa_2} - \frac{\dot{\kappa}_1}{\kappa_1} \right) \vec{u} \right)$ $\dot{\vec{N}}^{(3)} = \kappa_3 \vec{N}^{(2)} - \kappa_4 \vec{N}^{(4)}$
order 4	$u\ddot{\dot{u}} = 4\kappa_1 \ddot{\kappa}_1 + 3\dot{\kappa}_1^2 + \kappa_1^4 - \kappa_1^2 \kappa_2^2$ $\dot{u}\ddot{\dot{u}} = -6\kappa_1^3 \dot{\kappa}_1 + 3\kappa_1 \kappa_2^2 \dot{\kappa}_1 + 3\kappa_1^2 \kappa_2 \dot{\kappa}_2 - \kappa_1 \dot{\kappa}_1$ $\ddot{u}\ddot{\dot{u}} = -\dot{\kappa}_1 \ddot{\kappa}_1 + 4\kappa_1^3 \ddot{\kappa}_1 - 3\kappa_1 \kappa_2^2 \ddot{\kappa}_1 - \kappa_1^2 \kappa_2 \ddot{\kappa}_2 - 3\kappa_1^2 \dot{\kappa}_1^2 + 3\kappa_2^2 \dot{\kappa}_1^2 + \kappa_1^6 - 2\kappa_1^4 \kappa_2^2 + \kappa_1^2 \kappa_2^4 + \kappa_1^2 \kappa_2^2 \kappa_3^2$ $\dot{\dot{u}}\ddot{\dot{u}} = -\ddot{\kappa}_1 \dot{\kappa}_1 + 6\kappa_1^2 \dot{\kappa}_1 \dot{\kappa}_1 - \kappa_1 \kappa_2^2 \dot{\kappa}_3 \dot{\kappa}_1 - \kappa_1^2 \kappa_2 \kappa_3^2 \dot{\kappa}_2 - \kappa_1^2 \kappa_2^2 \kappa_3 \dot{\kappa}_3 - 3\kappa_2^2 \dot{\kappa}_1 \dot{\kappa}_1 + 9\kappa_1 \dot{\kappa}_1^3 + 4\kappa_1^3 \kappa_2^2 \dot{\kappa}_1 + 2\kappa_1^4 \kappa_2 \dot{\kappa}_2 - \kappa_1 \kappa_2^4 \dot{\kappa}_1 - 2\kappa_1^2 \kappa_2^3 \dot{\kappa}_2 + \kappa_1 \kappa_2^2 \dot{\kappa}_1 - 3\kappa_1^5 \dot{\kappa}_1 - \kappa_1^3 \dot{\kappa}_1 - 2\kappa_2 \dot{\kappa}_1 \kappa_1 \dot{\kappa}_2 - 6\kappa_2 \dot{\kappa}_1^2 \dot{\kappa}_2 - 3\kappa_1 \dot{\kappa}_1 \dot{\kappa}_2^2 - \kappa_1^2 \dot{\kappa}_2 \dot{\kappa}_2$ $\ddot{\dot{u}}^2 = -9\kappa_2^2 \kappa_3^2 \dot{\kappa}_1^2 - \kappa_1^2 \kappa_2^2 \kappa_3^2 \kappa_4^2 + 6\kappa_2^2 \dot{\kappa}_1 \dot{\kappa}_1 - 9\kappa_1^2 \kappa_2^2 \dot{\kappa}_2^2 - 14\kappa_1^3 \kappa_2^2 \dot{\kappa}_1 - 6\kappa_2^2 \kappa_3 \dot{\kappa}_1 \kappa_1 \dot{\kappa}_3 - 4\kappa_1^2 \kappa_3 \dot{\kappa}_2 \kappa_2 \dot{\kappa}_3 - 6\kappa_1 \dot{\kappa}_2 \kappa_1 \kappa_2 \kappa_3^2 + 30\kappa_1^2 \kappa_2^2 \dot{\kappa}_1^2 - \kappa_1^2 \kappa_2^2 \kappa_3^4 - 2\kappa_1^2 \kappa_2^4 \kappa_3^2 - \kappa_1^2 \kappa_2^2 \dot{\kappa}_3^2 + 2\kappa_1^4 \kappa_2^2 \kappa_3^2 - 3\kappa_1^6 \kappa_2^2 - 30\kappa_1^4 \dot{\kappa}_1^2 + 3\kappa_1^2 \kappa_2^4 - 9\kappa_2^4 \dot{\kappa}_1^2 + 30\kappa_1^3 \dot{\kappa}_1 \kappa_2 \dot{\kappa}_2 - 12\kappa_2^3 \dot{\kappa}_1 \kappa_1 \dot{\kappa}_2 + 8\kappa_1^5 \dot{\kappa}_1 + 16\kappa_1^2 \ddot{\kappa}_1^2 - 9\kappa_2^2 \ddot{\kappa}_1^2 - \kappa_1^2 \ddot{\kappa}_2^2 + 6\kappa_1 \kappa_2 \dot{\kappa}_2 \dot{\kappa}_1 - 18\kappa_1 \dot{\kappa}_2 \kappa_2 \ddot{\kappa}_1 - 6\kappa_1 \dot{\kappa}_2 \kappa_1 \ddot{\kappa}_2 + 6\kappa_1 \kappa_2^2 \kappa_3^2 \dot{\kappa}_1 + 2\kappa_1^2 \kappa_2 \kappa_3^2 \dot{\kappa}_2 - 6\kappa_2 \ddot{\kappa}_1 \kappa_1 \ddot{\kappa}_2 + 24\kappa_1 \ddot{\kappa}_1 \dot{\kappa}_1^2 - 12\kappa_1^2 \dot{\kappa}_1 \dot{\kappa}_1 - 2\kappa_1^4 \kappa_2 \ddot{\kappa}_2 + 6\kappa_1 \kappa_2^4 \ddot{\kappa}_1 + 2\kappa_1^2 \kappa_2^3 \ddot{\kappa}_2 - \dot{\kappa}_1^2$	$\vec{N}^{(4)} = \frac{1}{\kappa_1 \kappa_2 \kappa_3 \kappa_4} \left[-\ddot{\vec{u}} + \left(3\frac{\dot{\kappa}_1}{\kappa_1} + 2\frac{\dot{\kappa}_2}{\kappa_2} \frac{\dot{\kappa}_3}{\kappa_3} \right) \dot{\vec{u}} + \left(\kappa_1^2 - \kappa_2^2 - \kappa_3^2 + 3\frac{\dot{\kappa}_1}{\kappa_1} - 6\frac{\dot{\kappa}_1^2}{\kappa_1^2} - 2\frac{\dot{\kappa}_2^2}{\kappa_2^2} - 4\frac{\dot{\kappa}_1 \dot{\kappa}_2}{\kappa_1 \kappa_2} - 2\frac{\dot{\kappa}_1 \dot{\kappa}_3}{\kappa_1 \kappa_3} - \frac{\dot{\kappa}_2 \dot{\kappa}_3}{\kappa_2 \kappa_3} \right) \ddot{\vec{u}} + \left(\frac{\dot{\kappa}_1}{\kappa_1} + 6\frac{\dot{\kappa}_1^3}{\kappa_1^3} + (\kappa_1^2 + \kappa_2^2 + \kappa_3^2) \frac{\dot{\kappa}_1}{\kappa_1} - (2\kappa_1^2 + \kappa_2^2) \frac{\dot{\kappa}_2}{\kappa_2} - (\kappa_1^2 - \kappa_2^2) \frac{\dot{\kappa}_3}{\kappa_3} - 6\frac{\dot{\kappa}_1 \dot{\kappa}_1}{\kappa_1^2} - \frac{\dot{\kappa}_1 \dot{\kappa}_2}{\kappa_1 \kappa_2} - 2\frac{\dot{\kappa}_1 \dot{\kappa}_2}{\kappa_1 \kappa_2} - \frac{\dot{\kappa}_1 \dot{\kappa}_3}{\kappa_1 \kappa_3} + 4\frac{\dot{\kappa}_2 \dot{\kappa}_2}{\kappa_1^2 \kappa_2} + 2\frac{\dot{\kappa}_1 \dot{\kappa}_2^2}{\kappa_1 \kappa_2^2} + 2\frac{\dot{\kappa}_2 \kappa_3}{\kappa_1^2 \kappa_3} + \frac{\dot{\kappa}_1 \dot{\kappa}_2 \dot{\kappa}_3}{\kappa_1 \kappa_2 \kappa_3} \right) \dot{\vec{u}} + \kappa_1^2 \left(\frac{\dot{\kappa}_1}{\kappa_1} - \frac{\dot{\kappa}_2}{\kappa_2} + \kappa_3^2 + 2\frac{\dot{\kappa}_2^2}{\kappa_2^2} - 2\frac{\dot{\kappa}_1 \dot{\kappa}_2}{\kappa_1 \kappa_2} - \frac{\dot{\kappa}_1 \dot{\kappa}_3}{\kappa_1 \kappa_3} + \frac{\dot{\kappa}_2 \dot{\kappa}_3}{\kappa_2 \kappa_3} \right) \vec{u} \right]$ $\dot{\vec{N}}^{(4)} = \kappa_4 \vec{N}^{(3)} - \kappa_5 \vec{N}^{(5)}$

TABLE V:

d = 4

	Radiated momentum flux	Particular Projections
rank 0 tensor	$\frac{d\mathcal{P}_{4 0}^\mu}{d\tau} = \frac{4\pi}{3}\dot{u}^2 u^\mu$	$P_0^{(4 0)} = I_{4 0} = -\frac{4\pi}{3}\kappa_1^2$
vector	$\frac{d\mathcal{P}_{4 1}^\mu}{d\tau} = -\frac{8\pi}{3}\dot{u}^2 u^\mu$	$P_0^{(4 1)} = I_{4 1} = \frac{8\pi}{3}\kappa_1^2$
rank 2 tensor	$\frac{d\mathcal{P}_{4 2}^\mu}{d\tau} = -\frac{20\pi}{3}\dot{u}^2 u^\mu$	$P_0^{(4 2)} = I_{4 2} = \frac{20\pi}{3}\kappa_1^2$
gravity	$\frac{d\mathcal{P}_{4 g}^\mu}{d\tau} = \mathcal{P}_{4 2}^\mu - \frac{1}{2}\frac{d\mathcal{P}_{4 0}^\mu}{d\tau} = -\frac{22\pi}{3}\dot{u}^2 u^\mu$	$P_0^{(4 g)} = I_{4 g} = I_{4 2} = \frac{22\pi}{3}\kappa_1^2$
rank 3 tensor	$\frac{d\mathcal{P}_{4 3}^\mu}{d\tau} = -\frac{32\pi}{3}\dot{u}^2 u^\mu$	$P_0^{(4 3)} = I_{4 3} = \frac{32\pi}{3}\kappa_1^2$
rank s tensor	$\frac{d\mathcal{P}_{4 s}^\mu}{d\tau} = \frac{4-12s}{3}\pi\dot{u}^2 u^\mu$	$P_0^{(4 s)} = I_{4 s} = \frac{12s-4}{3}\pi\kappa_1^2$

$$\boxed{d = 6}$$

Radiated momentum flux	Particular Projections
$\frac{d\mathcal{P}_{6 0}^\mu}{d\tau} = -\frac{8\pi^2}{3}\dot{u}^4 u^\mu + \frac{8\pi^2}{15}\ddot{u}^2 u^\mu + \frac{32\pi^2}{35}(\dot{u}\ddot{u})\dot{u}^\mu - \frac{64\pi^2}{105}\dot{u}^2\ddot{u}^\mu$	$I_{6 0} = \frac{8\pi^2}{105}\left(20\kappa_1^4 + 7\dot{\kappa}_1^2 + 7\kappa_1^2\kappa_2^2\right)$
$\frac{d\mathcal{P}_{6 1}^\mu}{d\tau} = \frac{16\pi^2}{3}\dot{u}^4 u^\mu - \frac{32\pi^2}{15}\ddot{u}^2 u^\mu - \frac{16\pi^2}{7}(\dot{u}\ddot{u})\dot{u}^\mu + \frac{16\pi^2}{35}\dot{u}^2\ddot{u}^\mu$	$P_0^{(6 0)} = -\frac{8\pi^2}{15}(4\kappa_1^4 + \kappa_1^2\kappa_2^2)$ $I_{6 1} = \frac{32\pi^2}{105}\left(9\kappa_1^4 + 7\dot{\kappa}_1^2 + 7\kappa_1^2\kappa_2^2\right)$
$\frac{d\mathcal{P}_{6 2}^\mu}{d\tau} = \frac{16\pi^2}{3}\dot{u}^4 u^\mu - \frac{24\pi^2}{5}\ddot{u}^2 u^\mu - \frac{192\pi^2}{35}(\dot{u}\ddot{u})\dot{u}^\mu + \frac{32\pi^2}{21}\dot{u}^2\ddot{u}^\mu$	$P_0^{(6 1)} = \frac{16\pi^2}{15}(3\kappa_1^4 + 2\kappa_1^2\kappa_2^2)$ $I_{6 2} = \frac{8\pi^2}{105}\left(-13\kappa_1^4 + 63\dot{\kappa}_1^2 + 63\kappa_1^2\kappa_2^2\right)$
$\frac{d\mathcal{P}_{6 g}^\mu}{d\tau} = \frac{20\pi^2}{3}\dot{u}^4 u^\mu - \frac{76\pi^2}{15}\ddot{u}^2 u^\mu - \frac{208\pi^2}{35}(\dot{u}\ddot{u})\dot{u}^\mu + \frac{64}{35}\dot{u}^2\ddot{u}^\mu$	$P_0^{(6 2)} = \frac{8\pi^2}{15}(\kappa_1^4 + 9\kappa_1^2\kappa_2^2)$ $I_{6 g} = I_{6 2} - \frac{1}{2}I_{6 0} =$ $= \frac{4\pi^2}{105}\left(-46\kappa_1^4 + 119\dot{\kappa}_1^2 + 119\kappa_1^2\kappa_2^2\right)$
$\frac{d\mathcal{P}_{6 3}^\mu}{d\tau} = -\frac{8\pi^2}{3}\dot{u}^4 u^\mu - \frac{112\pi^2}{15}\ddot{u}^2 u^\mu - \frac{304\pi^2}{35}(\dot{u}\ddot{u})\dot{u}^\mu + \frac{272\pi^2}{105}\dot{u}^2\ddot{u}^\mu$	$P_0^{(6 g)} = \frac{4\pi^2}{15}(6\kappa_1^4 + 19\kappa_1^2\kappa_2^2)$ $I_{6 3} = \frac{8\pi^2}{105}\left(-167\kappa_1^4 + 98\dot{\kappa}_1^2 + 98\kappa_1^2\kappa_2^2\right)$
$\frac{d\mathcal{P}_{6 s}^\mu}{d\tau} = \pi^2 \left[\frac{19}{3} - (2s-3)^2\right] \dot{u}^4 u^\mu + \frac{8\pi^2}{15}(1-5s)\ddot{u}^2 u^\mu +$ $+ \frac{16\pi^2}{35}(2-7s)(\dot{u}\ddot{u})\dot{u}^\mu + \frac{16\pi^2}{105}(7s-4)\dot{u}^2\ddot{u}^\mu$	$P_0^{(6 3)} = -\frac{8\pi^2}{15}(19\kappa_1^4 - 14\kappa_1^2\kappa_2^2)$
	$I_{6 s} = -\pi^2 \left(4s^2 - \frac{124}{15}s + \frac{32}{21}\right) \kappa_1^4 +$ $+ \frac{8\pi^2}{15}(5s-1)\dot{\kappa}_1^2 + \frac{8\pi^2}{15}(5s-1)\kappa_1^2\kappa_2^2$ $P_0^{(6 s)} = -\pi^2 \left(4s^2 - \frac{28}{3}s + \frac{32}{15}\right) 4\kappa_1^4 +$ $+ \frac{8\pi^2}{15}(5s-1)\kappa_1^2\kappa_2^2$

$$\boxed{d = 8}$$

$\begin{aligned} \frac{d\mathcal{P}_{8 0}^\mu}{d\tau} &= \left\{ \frac{80\pi^3}{9} \dot{u}^6 - \frac{1312\pi^3}{189} (\dot{u}\ddot{u})^2 - \frac{320\pi^3}{189} \dot{u}^2 \ddot{u}^2 + \frac{352\pi^3}{189} \dot{u}^2 (\dot{u}\ddot{u}) + \frac{16\pi^3}{105} \dot{u}^2 \ddot{u}^2 \right\} u^\mu - \\ &\quad - \left\{ \frac{9056\pi^3}{2079} \dot{u}^2 (\dot{u}\ddot{u}) - \frac{64\pi^3}{189} \ddot{u}\ddot{u} \right\} \dot{u}^\mu + \left\{ \frac{4160\pi^3}{2079} \dot{u}^4 + \frac{64\pi^3}{189} \dot{u}\ddot{u} \right\} \ddot{u}^\mu - \frac{544\pi^3}{945} (\dot{u}\ddot{u}) \dot{u}^\mu \\ \frac{d\mathcal{P}_{8 1}^\mu}{d\tau} &= \left\{ -\frac{160\pi^3}{9} \dot{u}^6 + \frac{12448\pi^3}{945} (\dot{u}\ddot{u})^2 + \frac{5888\pi^3}{945} \dot{u}^2 \ddot{u}^2 - \frac{4864\pi^3}{945} \dot{u}^2 (\dot{u}\ddot{u}) - \frac{32\pi^3}{35} \dot{u}^2 \ddot{u}^2 \right\} u^\mu + \\ &\quad + \left\{ \frac{78272\pi^3}{10395} \dot{u}^2 (\dot{u}\ddot{u}) - \frac{1408\pi^3}{945} \ddot{u}\ddot{u} \right\} \dot{u}^\mu - \left\{ \frac{29888\pi^3}{10395} \dot{u}^4 + \frac{832\pi^3}{945} \dot{u}\ddot{u} \right\} \ddot{u}^\mu + \frac{64\pi^3}{189} (\dot{u}\ddot{u}) \dot{u}^\mu \\ \frac{d\mathcal{P}_{8 2}^\mu}{d\tau} &= \left\{ -\frac{5552\pi^3}{315} \dot{u}^6 + \frac{11296\pi^3}{945} (\dot{u}\ddot{u})^2 + \frac{2272\pi^3}{189} \dot{u}^2 \ddot{u}^2 - \frac{7456\pi^3}{945} \dot{u}^2 (\dot{u}\ddot{u}) - \frac{208\pi^3}{105} \dot{u}^2 \ddot{u}^2 \right\} u^\mu + \\ &\quad + \left\{ \frac{87776\pi^3}{10395} \dot{u}^2 (\dot{u}\ddot{u}) - \frac{448\pi^3}{135} \ddot{u}\ddot{u} \right\} \dot{u}^\mu - \left\{ \frac{55232\pi^3}{10395} \dot{u}^4 + \frac{1984\pi^3}{945} \dot{u}\ddot{u} \right\} \ddot{u}^\mu + \frac{1184\pi^3}{945} (\dot{u}\ddot{u}) \dot{u}^\mu \\ \frac{d\mathcal{P}_{8 g}^\mu}{d\tau} &= \left\{ -\frac{6952\pi^3}{315} \dot{u}^6 + \frac{14576\pi^3}{945} (\dot{u}\ddot{u})^2 + \frac{2432\pi^3}{189} \dot{u}^2 \ddot{u}^2 - \frac{8336\pi^3}{945} \dot{u}^2 (\dot{u}\ddot{u}) - \frac{72\pi^3}{35} \dot{u}^2 \ddot{u}^2 \right\} u^\mu + \\ &\quad + \left\{ \frac{110416\pi^3}{10395} \dot{u}^2 (\dot{u}\ddot{u}) - \frac{3296\pi^3}{945} \ddot{u}\ddot{u} \right\} \dot{u}^\mu - \left\{ \frac{9376\pi^3}{1485} \dot{u}^4 + \frac{2144\pi^3}{945} \dot{u}\ddot{u} \right\} \ddot{u}^\mu + \frac{208\pi^3}{135} (\dot{u}\ddot{u}) \dot{u}^\mu \\ \frac{d\mathcal{P}_{8 3}^\mu}{d\tau} &= \left\{ \frac{592\pi^3}{315} \dot{u}^6 - \frac{10016\pi^3}{945} (\dot{u}\ddot{u})^2 + \frac{14816\pi^3}{945} \dot{u}^2 \ddot{u}^2 - \frac{6016\pi^3}{945} \dot{u}^2 (\dot{u}\ddot{u}) - \frac{64\pi^3}{21} \dot{u}^2 \ddot{u}^2 \right\} u^\mu - \\ &\quad - \left\{ \frac{16768\pi^3}{10395} \dot{u}^2 (\dot{u}\ddot{u}) + \frac{4864\pi^3}{945} \ddot{u}\ddot{u} \right\} \dot{u}^\mu - \left\{ \frac{55232\pi^3}{10395} \dot{u}^4 + \frac{448\pi^3}{135} \dot{u}\ddot{u} \right\} \ddot{u}^\mu + \frac{2048\pi^3}{945} (\dot{u}\ddot{u}) \dot{u}^\mu \end{aligned}$	$\begin{aligned} P_0^{(8 0)} &= -\frac{16\pi^3}{315} (3\kappa_1^2 \kappa_2^2 \kappa_3^2 + 108\kappa_1^6 + \\ &\quad + 64\kappa_1^4 \kappa_2^2 + 3\kappa_1^2 \kappa_2^4) \\ P_0^{(8 1)} &= \frac{32\pi^3}{315} (9\kappa_1^2 \kappa_2^2 \kappa_3^2 + 72\kappa_1^6 + \\ &\quad + 94\kappa_1^4 \kappa_2^2 + 9\kappa_1^2 \kappa_2^4) \\ P_0^{(8 2)} &= \frac{16\pi^3}{315} (39\kappa_1^2 \kappa_2^2 \kappa_3^2 - 6\kappa_1^6 + \\ &\quad + 314\kappa_1^4 \kappa_2^2 + 39\kappa_1^2 \kappa_2^4) \\ P_0^{(8 g)} &= \frac{72\pi^3}{35} \kappa_1^2 \kappa_2^2 \kappa_3^2 + \frac{256\pi^3}{105} \kappa_1^6 + \\ &\quad + \frac{5536\pi^3}{315} \kappa_1^4 \kappa_2^2 + \frac{72\pi^3}{35} \kappa_1^2 \kappa_2^4 \\ P_0^{(8 3)} &= \frac{16\pi^3}{315} (60\kappa_1^2 \kappa_2^2 \kappa_3^2 - 411\kappa_1^6 + \\ &\quad + 314\kappa_1^4 \kappa_2^2 + 60\kappa_1^2 \kappa_2^4) \end{aligned}$
$\begin{aligned} \frac{d\mathcal{P}_{8 s}^\mu}{d\tau} &= \left\{ -\frac{16\pi^3}{9} \left(\frac{7s^3}{10} - \frac{135}{14} s^2 + \frac{838}{35} s - 5 \right) \dot{u}^6 + \frac{32\pi^3}{3} \left(-s^2 + \frac{101}{35} s - \frac{41}{63} \right) (\dot{u}\ddot{u})^2 + \right. \\ &\quad + \frac{16\pi^3}{3} \left(-\frac{s^2}{5} + \frac{59}{35} s - \frac{20}{63} \right) \dot{u}^2 \ddot{u}^2 + 32\pi^3 \left(\frac{s^2}{15} - \frac{2s}{7} + \frac{11}{189} \right) \dot{u}^2 (\dot{u}\ddot{u}) + \\ &\quad + \frac{16\pi^3}{15} \left(\frac{1}{7} - s \right) \dot{u}^2 \ddot{u}^2 \left. \right\} u^\mu + \left\{ \frac{32\pi^3}{7} \left(-\frac{6s^2}{5} + \frac{19}{5} s - \frac{283}{297} \right) \dot{u}^2 (\dot{u}\ddot{u}) + \frac{64\pi^3}{7} \left(\frac{1}{27} - \frac{s}{5} \right) \ddot{u}\ddot{u} \right\} \dot{u}^\mu + \\ &\quad + \left\{ \frac{64\pi^3}{21} \left(\frac{2s^2}{5} - 2s + \frac{65}{99} \right) \dot{u}^4 + \frac{64\pi^3}{21} \left(\frac{1}{9} - \frac{2s}{5} \right) \dot{u}\ddot{u} \right\} \ddot{u}^\mu + \frac{32\pi^3}{35} \left(s - \frac{17}{27} \right) (\dot{u}\ddot{u}) \dot{u}^\mu \end{aligned}$	$\begin{aligned} P_0^{(8 s)} &= \frac{16\pi^3}{105} (7s - 1) (\kappa_1^2 \kappa_2^2 \kappa_3^2 + \kappa_1^2 \kappa_2^4) + \\ &\quad + \frac{8\pi^3}{315} (49s^3 - 549s^2 + 1004s - 216) \kappa_1^6 - \\ &\quad - 16\pi^3 \left(\frac{s^2}{5} - s + \frac{64}{315} \right) \kappa_1^4 \kappa_2^2 \end{aligned}$

$$I_{8|0} = -\frac{16\pi^3}{1155} \left[276\kappa_1^6 + \frac{1892}{9} \kappa_1^4 \kappa_2^2 + \frac{1199}{3} \kappa_1^2 \dot{\kappa}_1^2 - 88\kappa_1^3 \ddot{\kappa}_1 + 11 \{ \kappa_1^2 \kappa_2^2 \kappa_3^2 + (\kappa_1 \kappa_2^2 - \ddot{\kappa}_1)^2 + (2\kappa_2 \dot{\kappa}_1 + \kappa_1 \dot{\kappa}_2)^2 \} \right]$$

$$I_{8|1} = \frac{32\pi^3}{385} \left[64\kappa_1^6 + \frac{2816}{27} \kappa_1^4 \kappa_2^2 + \frac{1100}{9} \kappa_1^2 \dot{\kappa}_1^2 - \frac{88}{3} \kappa_1^3 \ddot{\kappa}_1 + 11 \{ \kappa_1^2 \kappa_2^2 \kappa_3^2 + (\kappa_1 \kappa_2^2 - \ddot{\kappa}_1)^2 + (2\kappa_2 \dot{\kappa}_1 + \kappa_1 \dot{\kappa}_2)^2 \} \right]$$

$$I_{8|2} = \frac{16\pi^3}{1155} \left[-254\kappa_1^6 + \frac{8998}{9} \kappa_1^4 \kappa_2^2 + \frac{517}{3} \kappa_1^2 \dot{\kappa}_1^2 - 132\kappa_1^3 \ddot{\kappa}_1 + 143 \{ \kappa_1^2 \kappa_2^2 \kappa_3^2 + (\kappa_1 \kappa_2^2 - \ddot{\kappa}_1)^2 + (2\kappa_2 \dot{\kappa}_1 + \kappa_1 \dot{\kappa}_2)^2 \} \right]$$

$$I_{8|g} = \frac{8\pi^3}{1155} \left[-232\kappa_1^6 + \frac{19888}{9} \kappa_1^4 \kappa_2^2 + \frac{2233}{3} \kappa_1^2 \dot{\kappa}_1^2 - 352\kappa_1^3 \ddot{\kappa}_1 + 297 \{ \kappa_1^2 \kappa_2^2 \kappa_3^2 + (\kappa_1 \kappa_2^2 - \ddot{\kappa}_1)^2 + (2\kappa_2 \dot{\kappa}_1 + \kappa_1 \dot{\kappa}_2)^2 \} \right]$$

$$I_{8|3} = \frac{16\pi^3}{1155} \left[-1651\kappa_1^6 + \frac{8206}{9} \kappa_1^4 \kappa_2^2 - \frac{6248}{3} \kappa_1^2 \dot{\kappa}_1^2 + 220\kappa_1^3 \ddot{\kappa}_1 + 220 \{ \kappa_1^2 \kappa_2^2 \kappa_3^2 + (\kappa_1 \kappa_2^2 - \ddot{\kappa}_1)^2 + (2\kappa_2 \dot{\kappa}_1 + \kappa_1 \dot{\kappa}_2)^2 \} \right]$$

$$\begin{aligned} I_{8|s} &= \frac{16\pi^3}{105} \left[\left(\frac{49}{6} s^3 - \frac{167}{2} s^2 + \frac{406}{3} s - \frac{276}{11} \right) \kappa_1^6 - \left(21s^2 - 97s + \frac{172}{9} \right) \kappa_1^4 \kappa_2^2 - \left(77s^2 - 180s + \frac{109}{3} \right) \kappa_1^2 \dot{\kappa}_1^2 + \right. \\ &\quad \left. + 2(7s^2 - 9s + 4) \kappa_1^3 \ddot{\kappa}_1 + (7s - 1) \{ \kappa_1^2 \kappa_2^2 \kappa_3^2 + (\kappa_1 \kappa_2^2 - \ddot{\kappa}_1)^2 + (2\kappa_2 \dot{\kappa}_1 + \kappa_1 \dot{\kappa}_2)^2 \} \right] \end{aligned}$$

$$\boxed{d = 10}$$

$$\begin{aligned} \frac{d\mathcal{P}_{10|0}^\mu}{d\tau} = & \left\{ -\frac{1120\pi^4}{27}\dot{u}^8 - \frac{1088\pi^4}{315}(\dot{u}\ddot{u})^2 - \frac{8192\pi^4}{3465}\ddot{u}^4 + \frac{32\pi^4}{945}\ddot{u}^2 + \frac{28480\pi^4}{429}\dot{u}^2(\dot{u}\ddot{u})^2 + \frac{5344\pi^4}{1287}\dot{u}^4\ddot{u}^2 - \right. \\ & - \frac{8896\pi^4}{429}\dot{u}^4(\dot{u}\ddot{u}) - \frac{2176\pi^4}{495}\ddot{u}^2(\dot{u}\ddot{u}) - \frac{1280\pi^4}{693}(\ddot{u}\ddot{u})(\dot{u}\ddot{u}) - \frac{160\pi^4}{231}\dot{u}^2\ddot{u}^2 + \frac{164\pi^4}{77}\dot{u}^2(\ddot{u}\ddot{u}) + \frac{128\pi^4}{63}(\dot{u}\ddot{u})(\dot{u}\ddot{u}) \Big\} u^\mu + \\ & + \left\{ \frac{64\pi^4}{693}(\ddot{u}\ddot{u}) - \frac{38656\pi^4}{9009}(\ddot{u}\ddot{u})(\dot{u}\ddot{u}) + \frac{896\pi^4}{1287}\dot{u}^2(\dot{u}\ddot{u}) + \frac{2880\pi^4}{143}\dot{u}^4(\dot{u}\ddot{u}) - \frac{320\pi^4}{819}\dot{u}^2(\ddot{u}\ddot{u}) - \frac{18944\pi^4}{3003}\ddot{u}^2(\dot{u}\ddot{u}) \right\} \dot{u}^\mu - \\ & - \left\{ \frac{37760\pi^4}{3861}\dot{u}^6 + \frac{2560\pi^4}{1001}(\dot{u}\ddot{u})^2 + \frac{256\pi^4}{2079}(\ddot{u}\ddot{u}) - \frac{33472\pi^4}{9009}\dot{u}^2(\dot{u}\ddot{u}) - \frac{2624\pi^4}{9009}\dot{u}^2\ddot{u}^2 \right\} \ddot{u}^\mu + \\ & + \left\{ \frac{64\pi^4}{693}\ddot{u}\ddot{u} + \frac{33920\pi^4}{9009}\dot{u}^2(\dot{u}\ddot{u}) \right\} \ddot{u}^\mu - \left\{ \frac{1472\pi^4}{10395}\ddot{u}^2 - \frac{1856\pi^4}{10395}\dot{u}\ddot{u} - \frac{2176\pi^4}{9009}\dot{u}^4 \right\} \ddot{u}^\mu \end{aligned}$$

$$\begin{aligned} \frac{d\mathcal{P}_{10|1}^\mu}{d\tau} = & \left\{ \frac{2240\pi^4}{27}\dot{u}^8 + \frac{704\pi^4}{105}(\dot{u}\ddot{u})^2 + \frac{7552\pi^4}{1485}\ddot{u}^4 - \frac{256\pi^4}{945}\ddot{u}^2 - \frac{15680\pi^4}{117}\dot{u}^2(\dot{u}\ddot{u})^2 - \frac{320\pi^4}{13}\dot{u}^4\ddot{u}^2 + \right. \\ & + \frac{5440\pi^4}{17}\dot{u}^4(\dot{u}\ddot{u}) + \frac{63424\pi^4}{10395}\ddot{u}^2(\dot{u}\ddot{u}) + \frac{640\pi^4}{77}(\dot{u}\ddot{u})(\ddot{u}\ddot{u}) + \frac{7360\pi^4}{2079}\dot{u}^2\ddot{u}^2 - \frac{7424\pi^4}{2079}\dot{u}^2(\ddot{u}\ddot{u}) - \frac{128\pi^4}{21}(\dot{u}\ddot{u})(\dot{u}\ddot{u}) \Big\} u^\mu - \\ & - \left\{ \frac{1216\pi^4}{2079}(\dot{u}\ddot{u}) + \frac{4352\pi^4}{819}(\dot{u}\ddot{u})(\dot{u}\ddot{u}) - \frac{1984\pi^4}{1001}\dot{u}^2(\ddot{u}\ddot{u}) - \frac{50560\pi^4}{1287}\dot{u}^4(\dot{u}\ddot{u}) + \frac{8000\pi^4}{3861}\dot{u}^2(\ddot{u}\ddot{u}) + \frac{112064\pi^4}{9009}\ddot{u}^2(\dot{u}\ddot{u}) \right\} \dot{u}^\mu + \\ & + \left\{ \frac{59200\pi^4}{3861}\dot{u}^6 - \frac{3200\pi^4}{819}(\dot{u}\ddot{u})^2 - \frac{128\pi^4}{231}(\ddot{u}\ddot{u}) + \frac{58048\pi^4}{9009}\dot{u}^2(\dot{u}\ddot{u}) - \frac{53632\pi^4}{27027}\dot{u}^2\ddot{u}^2 \right\} \ddot{u}^\mu - \\ & - \left\{ \frac{512\pi^4}{2079}\ddot{u}\ddot{u} - \frac{139520\pi^4}{27027}\dot{u}^2(\dot{u}\ddot{u}) \right\} \ddot{u}^\mu + \left\{ \frac{128\pi^4}{2079}\ddot{u}^2 + \frac{64\pi^4}{693}\dot{u}\ddot{u} + \frac{4288\pi^4}{27027}\dot{u}^4 \right\} \ddot{u}^\mu \end{aligned}$$

$$\begin{aligned} \frac{d\mathcal{P}_{10|2}^\mu}{d\tau} = & \left\{ \frac{160000\pi^4}{2079}\dot{u}^8 + \frac{2048\pi^4}{315}(\dot{u}\ddot{u})^2 + \frac{10016\pi^4}{1485}\ddot{u}^4 - \frac{544\pi^4}{945}\ddot{u}^2 - \frac{1429504\pi^4}{9009}\dot{u}^2(\dot{u}\ddot{u})^2 - \frac{1424480\pi^4}{27027}\dot{u}^4\ddot{u}^2 + \right. \\ & + \frac{210304\pi^4}{3861}\dot{u}^4(\dot{u}\ddot{u}) + \frac{4096\pi^4}{2079}\ddot{u}^2(\dot{u}\ddot{u}) + \frac{59776\pi^4}{3465}(\dot{u}\ddot{u})(\ddot{u}\ddot{u}) + \frac{74464\pi^4}{10395}\dot{u}^2\ddot{u}^2 - \frac{70208\pi^4}{10395}\dot{u}^2(\ddot{u}\ddot{u}) - \frac{3328\pi^4}{315}(\dot{u}\ddot{u})(\dot{u}\ddot{u}) \Big\} u^\mu - \\ & - \left\{ \frac{2624\pi^4}{2079}(\dot{u}\ddot{u}) + \frac{37120\pi^4}{27027}(\dot{u}\ddot{u})(\dot{u}\ddot{u}) - \frac{89344\pi^4}{27027}\dot{u}^2(\dot{u}\ddot{u}) - \frac{469120\pi^4}{9009}\dot{u}^4(\dot{u}\ddot{u}) + \frac{122560\pi^4}{27027}\dot{u}^2(\ddot{u}\ddot{u}) + \frac{476800\pi^4}{27027}\ddot{u}^2(\dot{u}\ddot{u}) \right\} \dot{u}^\mu + \\ & + \left\{ \frac{49984\pi^4}{2457}\dot{u}^6 - \frac{188800\pi^4}{27027}(\dot{u}\ddot{u})^2 - \frac{2560\pi^4}{2079}(\ddot{u}\ddot{u}) + \frac{37952\pi^4}{3861}\dot{u}^2(\dot{u}\ddot{u}) - \frac{57536\pi^4}{9009}\dot{u}^2\ddot{u}^2 \right\} \ddot{u}^\mu - \\ & - \left\{ \frac{1216\pi^4}{2079}\ddot{u}\ddot{u} - \frac{11392\pi^4}{1001}\dot{u}^2(\dot{u}\ddot{u}) \right\} \ddot{u}^\mu + \left\{ \frac{2752\pi^4}{10395}\ddot{u}^2 + \frac{3776\pi^4}{10395}\dot{u}\ddot{u} + \frac{66368\pi^4}{135135}\dot{u}^4 \right\} \ddot{u}^\mu \end{aligned}$$

$$\begin{aligned}
\frac{d\mathcal{P}_{10|g}^\mu}{d\tau} = & \left\{ \frac{203120\pi^4}{2079} \dot{u}^8 + \frac{288\pi^4}{35} (\dot{u}\ddot{u})^2 + \frac{16480\pi^4}{2079} \ddot{u}^4 - \frac{16\pi^4}{27} \ddot{u}^2 - \frac{1728544\pi^4}{9009} \dot{u}^2 (\ddot{u})^2 - \frac{1480592\pi^4}{27027} \dot{u}^4 \ddot{u}^2 + \right. \\
& + \frac{250336\pi^4}{3861} \dot{u}^4 (\dot{u}\ddot{u}) + \frac{43328\pi^4}{10395} \ddot{u}^2 (\dot{u}\ddot{u}) + \frac{20992\pi^4}{1155} (\ddot{u}\ddot{u})(\ddot{u}\ddot{u}) + \frac{11152\pi^4}{1485} \dot{u}^2 \ddot{u}^2 - \frac{74528\pi^4}{10395} \dot{u}^2 (\ddot{u}\ddot{u}) - \frac{1216\pi^4}{105} (\ddot{u}\ddot{u})(\ddot{u}\ddot{u}) \left. \right\} u^\mu - \\
& - \left\{ \frac{2720\pi^4}{2079} (\ddot{u}\ddot{u}) + \frac{95104\pi^4}{27027} (\ddot{u}\ddot{u})(\dot{u}\ddot{u}) - \frac{98752\pi^4}{27027} \dot{u}^2 (\ddot{u}\ddot{u}) - \frac{559840\pi^4}{9009} \dot{u}^4 (\dot{u}\ddot{u}) + \frac{127840\pi^4}{27027} \dot{u}^2 (\ddot{u}\ddot{u}) + \frac{562048\pi^4}{27027} \ddot{u}^2 (\dot{u}\ddot{u}) \right\} \dot{u}^\mu + \\
& + \left\{ \frac{75776\pi^4}{3003} \dot{u}^6 - \frac{223360\pi^4}{27027} (\ddot{u}\ddot{u})^2 - \frac{128\pi^4}{99} (\ddot{u}\ddot{u}) + \frac{315872\pi^4}{27027} \dot{u}^2 (\dot{u}\ddot{u}) - \frac{8032\pi^4}{1287} \dot{u}^2 \ddot{u}^2 \right\} \ddot{u}^\mu - \\
& - \left\{ \frac{1312\pi^4}{2079} \dot{u}\ddot{u} - \frac{119488\pi^4}{9009} \dot{u}^2 (\ddot{u}\ddot{u}) \right\} \dot{u}^\mu + \left\{ \frac{3488\pi^4}{10395} \ddot{u}^2 + \frac{224\pi^4}{495} \dot{u}\ddot{u} + \frac{82688\pi^4}{135135} \dot{u}^4 \right\} \ddot{u}^\mu
\end{aligned}$$

$$\begin{aligned}
\frac{d\mathcal{P}_{10|3}^\mu}{d\tau} = & \left\{ -\frac{3104\pi^4}{231} \dot{u}^8 - \frac{256\pi^4}{63} (\dot{u}\ddot{u})^2 + \frac{9056\pi^4}{3465} \ddot{u}^4 - \frac{832\pi^4}{945} \ddot{u}^2 - \frac{105152\pi^4}{3003} \dot{u}^2 (\ddot{u})^2 - \frac{3485408\pi^4}{45045} \dot{u}^4 \ddot{u}^2 + \right. \\
& + \frac{637568\pi^4}{45045} \dot{u}^4 (\dot{u}\ddot{u}) - \frac{6464\pi^4}{385} \ddot{u}^2 (\dot{u}\ddot{u}) + \frac{86528\pi^4}{3465} (\ddot{u}\ddot{u})(\ddot{u}\ddot{u}) + \frac{35264\pi^4}{3465} \dot{u}^2 \ddot{u}^2 - \frac{30208\pi^4}{3465} \dot{u}^2 (\ddot{u}\ddot{u}) - \frac{512\pi^4}{45} (\ddot{u}\ddot{u})(\dot{u}\ddot{u}) \left. \right\} u^\mu - \\
& - \left\{ \frac{64\pi^4}{33} (\ddot{u}\ddot{u}) - \frac{2304\pi^4}{143} (\ddot{u}\ddot{u})(\dot{u}\ddot{u}) - \frac{29504\pi^4}{9009} \dot{u}^2 (\ddot{u}\ddot{u}) - \frac{237376\pi^4}{9009} \dot{u}^4 (\dot{u}\ddot{u}) + \frac{63040\pi^4}{9009} \dot{u}^2 (\ddot{u}\ddot{u}) + \frac{1088\pi^4}{117} \ddot{u}^2 (\dot{u}\ddot{u}) \right\} \dot{u}^\mu + \\
& + \left\{ \frac{35968\pi^4}{3861} \dot{u}^6 - \frac{60160\pi^4}{9009} (\ddot{u}\ddot{u})^2 - \frac{3968\pi^4}{2079} (\ddot{u}\ddot{u}) + \frac{58048\pi^4}{9009} \dot{u}^2 (\dot{u}\ddot{u}) - \frac{121600\pi^4}{9009} \dot{u}^2 \ddot{u}^2 \right\} \ddot{u}^\mu - \\
& - \left\{ \frac{640\pi^4}{693} \dot{u}\ddot{u} - \frac{134144\pi^4}{9009} \dot{u}^2 (\ddot{u}\ddot{u}) \right\} \dot{u}^\mu + \left\{ \frac{4864\pi^4}{10395} \ddot{u}^2 + \frac{6592\pi^4}{10395} \dot{u}\ddot{u} + \frac{4864\pi^4}{6435} \dot{u}^4 \right\} \ddot{u}^\mu
\end{aligned}$$

$$P_0^{(10|0)} = -\frac{32\pi^4}{135135} [82368\kappa_1^8 + 80652\kappa_1^6\kappa_2^2 + 12012\kappa_1^4\kappa_2^4 + 6149\kappa_1^4\kappa_2^2\kappa_3^2 + 143(\kappa_1^2\kappa_2^6 + \kappa_1^2\kappa_2^2\kappa_3^2\kappa_4^2 + 2\kappa_1^2\kappa_2^4\kappa_3^2 + \kappa_1^2\kappa_2^2\kappa_3^4)]$$

$$P_0^{(10|1)} = \frac{64\pi^4}{135135} [51480\kappa_1^8 + 97812\kappa_1^6\kappa_2^2 + 25311\kappa_1^4\kappa_2^4 + 13871\kappa_1^4\kappa_2^2\kappa_3^2 + 572(\kappa_1^2\kappa_2^6 + \kappa_1^2\kappa_2^2\kappa_3^2\kappa_4^2 + 2\kappa_1^2\kappa_2^4\kappa_3^2 + \kappa_1^2\kappa_2^2\kappa_3^4)]$$

$$P_0^{(10|2)} = \frac{32\pi^4}{135135} [-23634\kappa_1^8 + 247104\kappa_1^6\kappa_2^2 + 99099\kappa_1^4\kappa_2^4 + 53911\kappa_1^4\kappa_2^2\kappa_3^2 + 2431(\kappa_1^2\kappa_2^6 + \kappa_1^2\kappa_2^2\kappa_3^2\kappa_4^2 + 2\kappa_1^2\kappa_2^4\kappa_3^2 + \kappa_1^2\kappa_2^2\kappa_3^4)]$$

$$P_0^{(10|g)} = \frac{16\pi^4}{135135} [35100\kappa_1^8 + 574860\kappa_1^6\kappa_2^2 + 210210\kappa_1^4\kappa_2^4 + 113971\kappa_1^4\kappa_2^2\kappa_3^2 + 5005(\kappa_1^2\kappa_2^6 + \kappa_1^2\kappa_2^2\kappa_3^2\kappa_4^2 + 2\kappa_1^2\kappa_2^4\kappa_3^2 + \kappa_1^2\kappa_2^2\kappa_3^4)]$$

$$P_0^{(10|3)} = \frac{32\pi^4}{135135} [-302419\kappa_1^8 + 108537\kappa_1^6\kappa_2^2 + 133419\kappa_1^4\kappa_2^4 + 72358\kappa_1^4\kappa_2^2\kappa_3^2 + 3718(\kappa_1^2\kappa_2^6 + \kappa_1^2\kappa_2^2\kappa_3^2\kappa_4^2 + 2\kappa_1^2\kappa_2^4\kappa_3^2 + \kappa_1^2\kappa_2^2\kappa_3^4)]$$

$$\begin{aligned}
I_{10|0} = & -\frac{32\pi^4}{135135} (66524\kappa_1^6\kappa_2^2 + 150140\kappa_1^4\kappa_1^2 + 11336\kappa_1^4\kappa_2^4 - 2704\kappa_1^4\kappa_2\ddot{\kappa}_2 - 22048\kappa_1^3\ddot{\kappa}_1\kappa_2^2 + 18278\kappa_1\ddot{\kappa}_1\dot{\kappa}_1^2 - 8684\kappa_1^2\dot{\kappa}_1\dot{\kappa}_1 + \\
& + 6903\dot{\kappa}_1^4 + 5629\kappa_1^4\kappa_2^2\kappa_3^2 + 55872\kappa_1^8 - 41904\kappa_1^5\ddot{\kappa}_1 + 12207\kappa_1^2\ddot{\kappa}_1^2 + 37440\kappa_1^3\dot{\kappa}_1\kappa_2\dot{\kappa}_2 + 2925\kappa_1^4\dot{\kappa}_2^2 + 48880\kappa_1^2\dot{\kappa}_1^2\kappa_2^2 + \\
& + 143[(\dot{\kappa}_1 - 3\dot{\kappa}_1\kappa_2^2 - 3\kappa_1\kappa_2\dot{\kappa}_2)^2 + (3\ddot{\kappa}_1\kappa_2 + 3\dot{\kappa}_1\dot{\kappa}_2 + \kappa_1\ddot{\kappa}_2 - \kappa_1\kappa_2^3)^2 + (2\kappa_1\dot{\kappa}_2\kappa_3 + \kappa_1\kappa_2\dot{\kappa}_3 + 3\dot{\kappa}_1\kappa_2\kappa_3)^2 + \\
& + \kappa_1^2\kappa_2^2\kappa_3^2(2\kappa_2^2 + \kappa_3^2 + \kappa_4^2) - 2\kappa_1\kappa_2\kappa_3^2(3\ddot{\kappa}_1\kappa_2 + \kappa_1\ddot{\kappa}_2 + 3\dot{\kappa}_1\dot{\kappa}_2)])
\end{aligned}$$

$$\begin{aligned}
I_{10|1} = & \frac{64\pi^4}{135135} (82152\kappa_1^6\kappa_2^2 + 109712\kappa_1^4\kappa_1^2 + 24076\kappa_1^4\kappa_2^4 - 5226\kappa_1^4\kappa_2\ddot{\kappa}_2 - 41028\kappa_1^3\ddot{\kappa}_1\kappa_2^2 + 15600\kappa_1\ddot{\kappa}_1\dot{\kappa}_1^2 - 10816\kappa_1^2\dot{\kappa}_1\dot{\kappa}_1 + \\
& + 5200\dot{\kappa}_1^4 + 12701\kappa_1^4\kappa_2^2\kappa_3^2 + 36000\kappa_1^8 - 32400\kappa_1^5\ddot{\kappa}_1 + 11700\kappa_1^2\ddot{\kappa}_1^2 + 64220\kappa_1^3\dot{\kappa}_1\kappa_2\dot{\kappa}_2 + 7475\kappa_1^4\dot{\kappa}_2^2 + 92248\kappa_1^2\dot{\kappa}_1^2\kappa_2^2 + \\
& + 572[(\dot{\kappa}_1 - 3\dot{\kappa}_1\kappa_2^2 - 3\kappa_1\kappa_2\dot{\kappa}_2)^2 + (3\ddot{\kappa}_1\kappa_2 + 3\dot{\kappa}_1\dot{\kappa}_2 + \kappa_1\ddot{\kappa}_2 - \kappa_1\kappa_2^3)^2 + (2\kappa_1\dot{\kappa}_2\kappa_3 + \kappa_1\kappa_2\dot{\kappa}_3 + 3\dot{\kappa}_1\kappa_2\kappa_3)^2 + \\
& + \kappa_1^2\kappa_2^2\kappa_3^2(2\kappa_2^2 + \kappa_3^2 + \kappa_4^2) - 2\kappa_1\kappa_2\kappa_3^2(3\ddot{\kappa}_1\kappa_2 + \kappa_1\ddot{\kappa}_2 + 3\dot{\kappa}_1\dot{\kappa}_2)])
\end{aligned}$$

$$\begin{aligned}
I_{10|2} = & \frac{32\pi^4}{135135} (187782\kappa_1^6\kappa_2^2 + 8980\kappa_1^4\kappa_1^2 + 93483\kappa_1^4\kappa_2^4 - 18460\kappa_1^4\kappa_2\ddot{\kappa}_2 - 139828\kappa_1^3\ddot{\kappa}_1\kappa_2^2 + 13754\kappa_1\ddot{\kappa}_1\dot{\kappa}_1^2 - 31356\kappa_1^2\dot{\kappa}_1\dot{\kappa}_1 + \\
& + 3250\dot{\kappa}_1^4 + 48711\kappa_1^4\kappa_2^2\kappa_3^2 - 44606\kappa_1^8 - 2478\kappa_1^5\ddot{\kappa}_1 + 12675\kappa_1^2\ddot{\kappa}_1^2 + 232544\kappa_1^3\dot{\kappa}_1\kappa_2\dot{\kappa}_2 + 30251\kappa_1^4\dot{\kappa}_2^2 + 353522\kappa_1^2\dot{\kappa}_1^2\kappa_2^2 + \\
& + 2431[(\dot{\kappa}_1 - 3\dot{\kappa}_1\kappa_2^2 - 3\kappa_1\kappa_2\dot{\kappa}_2)^2 + (3\ddot{\kappa}_1\kappa_2 + 3\dot{\kappa}_1\dot{\kappa}_2 + \kappa_1\ddot{\kappa}_2 - \kappa_1\kappa_2^3)^2 + (2\kappa_1\dot{\kappa}_2\kappa_3 + \kappa_1\kappa_2\dot{\kappa}_3 + 3\dot{\kappa}_1\kappa_2\kappa_3)^2 + \\
& + \kappa_1^2\kappa_2^2\kappa_3^2(2\kappa_2^2 + \kappa_3^2 + \kappa_4^2) - 2\kappa_1\kappa_2\kappa_3^2(3\ddot{\kappa}_1\kappa_2 + \kappa_1\ddot{\kappa}_2 + 3\dot{\kappa}_1\dot{\kappa}_2)])
\end{aligned}$$

$$\begin{aligned}
I_{10|g} = & \frac{32\pi^4}{135135} (221044\kappa_1^6\kappa_2^2 + 84050\kappa_1^4\kappa_1^2 + 99151\kappa_1^4\kappa_2^4 - 19812\kappa_1^4\kappa_2\ddot{\kappa}_2 - 150852\kappa_1^3\ddot{\kappa}_1\kappa_2^2 + 22893\kappa_1\ddot{\kappa}_1\dot{\kappa}_1^2 - 35698\kappa_1^2\dot{\kappa}_1\dot{\kappa}_1 + \\
& + \frac{13403}{2}\dot{\kappa}_1^4 + \frac{103051}{2}\kappa_1^4\kappa_2^2\kappa_3^2 - 16670\kappa_1^8 - 23430\kappa_1^5\ddot{\kappa}_1 + \frac{37557}{2}\kappa_1^2\ddot{\kappa}_1^2 + 251264\kappa_1^3\dot{\kappa}_1\kappa_2\dot{\kappa}_2 + \frac{63427}{2}\kappa_1^4\dot{\kappa}_2^2 + 377962\kappa_1^2\dot{\kappa}_1^2\kappa_2^2 + \\
& + 5005[(\dot{\kappa}_1 - 3\dot{\kappa}_1\kappa_2^2 - 3\kappa_1\kappa_2\dot{\kappa}_2)^2 + (3\ddot{\kappa}_1\kappa_2 + 3\dot{\kappa}_1\dot{\kappa}_2 + \kappa_1\ddot{\kappa}_2 - \kappa_1\kappa_2^3)^2 + (2\kappa_1\dot{\kappa}_2\kappa_3 + \kappa_1\kappa_2\dot{\kappa}_3 + 3\dot{\kappa}_1\kappa_2\kappa_3)^2 + \\
& + \kappa_1^2\kappa_2^2\kappa_3^2(2\kappa_2^2 + \kappa_3^2 + \kappa_4^2) - 2\kappa_1\kappa_2\kappa_3^2(3\ddot{\kappa}_1\kappa_2 + \kappa_1\ddot{\kappa}_2 + 3\dot{\kappa}_1\dot{\kappa}_2)])
\end{aligned}$$

$$\begin{aligned}
I_{10|3} = & \frac{32\pi^4}{135135} (38659\kappa_1^6\kappa_2^2 - 677225\kappa_1^4\dot{\kappa}_1^2 + 124657\kappa_1^4\kappa_2^4 - 21320\kappa_1^4\kappa_2\ddot{\kappa}_2 - 151268\kappa_1^3\ddot{\kappa}_1\kappa_2^2 - 70616\kappa_1\ddot{\kappa}_1\dot{\kappa}_1^2 - 20488\kappa_1^2\dot{\kappa}_1\dot{\kappa}_1 - \\
& - 28353\dot{\kappa}_1^4 + 64298\kappa_1^4\kappa_2^2\kappa_3^2 - 263119\kappa_1^8 + 182538\kappa_1^5\ddot{\kappa}_1 - 44382\kappa_1^2\ddot{\kappa}_1^2 + 274872\kappa_1^3\dot{\kappa}_1\kappa_2\dot{\kappa}_2 + 42978\kappa_1^4\dot{\kappa}_2^2 + 458198\kappa_1^2\dot{\kappa}_1^2\kappa_2^2 + \\
& + 3718[(\dot{\kappa}_1 - 3\dot{\kappa}_1\kappa_2^2 - 3\kappa_1\kappa_2\dot{\kappa}_2)^2 + (3\ddot{\kappa}_1\kappa_2 + 3\dot{\kappa}_1\dot{\kappa}_2 + \kappa_1\ddot{\kappa}_2 - \kappa_1\kappa_2^3)^2 + (2\kappa_1\dot{\kappa}_2\kappa_3 + \kappa_1\kappa_2\dot{\kappa}_3 + 3\dot{\kappa}_1\kappa_2\kappa_3)^2 + \\
& + \kappa_1^2\kappa_2^2\kappa_3^2(2\kappa_2^2 + \kappa_3^2 + \kappa_4^2) - 2\kappa_1\kappa_2\kappa_3^2(3\ddot{\kappa}_1\kappa_2 + \kappa_1\ddot{\kappa}_2 + 3\dot{\kappa}_1\dot{\kappa}_2)])
\end{aligned}$$

Appendix: MAPLE programs

This Appendix contains a MAPLE program that allows one to obtain the quantities listed in the tables with the help of a computer.

MAPLE program

The MAPLE program we used in calculations looks as follows (note that $uu[i, j]$ in the program is nothing but U_{ij} , $A[p, q]$ is $A_q^p U^{2p-q+1}$, $k[i](z)$ is $\kappa_i(\tau)$ etc.; $P[d, m]$'s are the angular integrals, formula (16)):

```
> #input I -> L then input P and apply L to P then input uu(k) and convert into k
>
> # u[i] = d^iu: uu[i,j] = (d^iu d^ju):
>
>
> M:=4: J:=3:
> A:=array(0..M,0..M): B:=array(0..M,0..M): II:=array(0..J,0..2*M+2): GG:=array(0..M,0..M):
> k:=array(0..M): K:=array(0..M): N:=array(0..M): tu:=array(0..M): u:= array(0..M):
> uu:=array(symmetric,0..M,0..M): v:=array(0..M): VV:=array(0..M,0..M):
> for i from 0 to M do: unassign('u[i]', 'uu[i,i]', 'tu[i]'):
> for j from 0 to i-1 do:
> unassign('uu[j,i]', 'VV[j,i]'):
> od: od:
>
> #----- Parameters A[p,q] -----
> VV[0,0]:=1/V(y):
> for p from 0 to M-1 do:
> VV[p+1,0]:=diff(VV[p,0],y)/V(y):
> for q from 0 to p do:
> VV[p+1,q+1]:=(diff(VV[p,q+1],y) + VV[p,q])/V(y):
> od: od:
> V(y):=add(v[n]*y^n/n!,n=0..M):
> for p from 0 to M do: for q from 0 to M do:
> A[p,q]:=simplify(v[0]^(2*p+1-q)*eval(VV[p,q],y=0)):
> od:od:
>
> #----- list entries of TABLE II-----
> for p to M do
> for j from 0 to p do
> print(A[p,j]):
> od: od:
>
>
> #---entries of TABLE III: Polynomials P[d,m]-----
>
> x:=t*t + a[0]*u+add(a[l1]*u[l1],l1=1..M): xu:= t*tu[0]+add(a[l2]*uu[0,l2],l2=0..M):
> x2:=t^2*t2 + a[0]^2 + add(a[n1]*a[n1]*uu[n1,n1],n1=1..M)+add(2*t*a[l3]*tu[l3],l3=0..M)+
> 2*add(add(a[n]*a[l4]*uu[n,l4],n=0..l4-1),l4=1..M): xL:= t*Lt+a[0]*Lu+add(a[l5]*Lu[l5],l5=1..M):
>
> x; xu; x2; xL; #----- print for illustration purposes -----
>
>
> for p to M do
> for l to p do
> P[2*p+2,2*1]:=2^(p+1)*Pi^p*2^(p-1+1)*(p-1+1)!/(2*p-1+2*1)!*add((-1)^n*(2*1)!*
> (xu)^(2*1-2*n)*(x2)^n/2^n/n!/(2*1-2*n)!*2^(p-1+2*1-n)*(p-1+2*1-n)!/2^(p-1+1)/(p-1+1)!,n=0..1):
> od: od:
```

```

>
> for p to M do
> for l from 0 to p do
> P[2*p+2,2*l+1]:=2^(p+1)*Pi^p*2^(p-1+1)*(p-1+1)!/(2*p-1+2*1)!*add((-1)^n*(2*1+1)!*
> (xu)^(2*1+1-2*n)*(x2)^n/2^n/n!/(2*1+1-2*n)!*2^(1-n)*(p+2*1-n)!/(p+1)!,n=0..1):
> od: od:
>
> #-----G[d,q,r]-----
>
> #-----expect long calculation-----
> for p to M do
>   for q from 0 to p do
>     for r from 0 to p do
> JJ:=A[p,q]*A[p,r]*P[2*p+2,2*p+1-q-r]:
>     for j from 0 to M do
> LLL:=add(ii!*coeff(coeff(JJ,v[j],ii),a[j],ii),ii=0..2*M):
>     JJ:=LLL:
>   od:
> G[2*p+2,q,r]:=diff(LLL,t)/(2*p+1-q-r)!:
>   od: od:
> od:
> #-----expect long calculation-----
>
> #---relations between uu-----
>
>   uu[0,0]:=1:
>   for j to M do
>     uu:=add(j!*uu[n,j-n]/n!/(j-n)!,n=1..j-1):uu[0,j]:=-uuu/2:
>     uu[0,j]:=-uuu/2:
>   od:
>
> #-----entries of TABLE V: intensities of radiation of spin-J particles
>
> for p to M do
>   for q from 0 to p do
>     for r from 0 to p do
> GG[q,r]:=simplify(G[2*p+2,q,r]): #print(simplify(G[2*p+2,q,r]),2*p+2,q,r,GG[q,r]):
>   od: od:
> II[0,2*p+2]:=GG[0,0]:
>   for j to J do
>     for q from 0 to p do
>       for r from 0 to p do
> GG[q,r]:= add(add( uu[q1,r1]*GG[q+q1,r+r1],q1=0..p-q),r1=0..p-r):
>       unassign('q1','r1'):
>     od: od:
>   II[j,2*p+2]:=simplify(GG[0,0]):
>   od:
> od:
>
> for p to M do
> for j from 0 to J do
> print(2*p+2,j,II[j,2*p+2]):
> od: od:
>
> #-----FRENET CALCULATION (TABLE IV)-----
> #-----expect long calculation-----
> uu[0,0]:=1:uu[1,0]:=0:uu[1,1]:=-k[1](z)^2: N[0]:=u[0]:N[1]:=1/k[1](z)*u[1]:
> k[0](z):=1: K[0](z):=1:K[1](z):=k[1](z):

```

```

> for p from 1 to M-1 do:
> uu[p,p+1]:=diff(uu[p,p],z)/2:
> K[p+1](z):=-K[p](z)*k[p+1](z):
> for i from 0 to p-1 do:
> uu[i,p+1]:=-uu[i+1,p]+diff(uu[i,p],z)
> od:
> B[p+1,0]:=uu[0,p+1]:
> for i from 1 to p do
> B[p+1,i]:=-add(uu[jj,p+1]*diff(N[i],u[jj]),jj=0..p):
> od:
> N[p+1]:=(u[p+1]-add(B[p+1,ii]*N[ii],ii=0..p))/K[p+1](z):
> uu[p+1,p+1]:=uu[0,p+1]^2 - K[p+1](z)^2 -add(B[p+1,ii]^2,ii=1..p):od:
> #-----expect long calculation-----
>
> #----list entries of TABLE IV-----
> for p from 0 to M do
> for j from 0 to p do uu[p,j]:=simplify(uu[p,j]):
> print(uu[p,j]):
> od: NN[p]:=simplify(N[p]):print(p,NN[p]): od:
> #-----end of FRENET-----
>
> #-----intensities through FRENET (TABLE V)-----
> for p to M do
> for j from 0 to J do
> III[j,2*p+2]:=simplify(II[j,2*p+2]):
> print(2*p+2,j,III[j,2*p+2]):
> od: od:
> #-----P_0 -- projection to u (TABLE V)-----
> for p to M do
> for j from 0 to J do
> IIP[j,2*p+2]:=subs(tu[1]=0,tu[2]=0,tu[3]=0,tu[4]=0,III[j,2*p+2]):
> print(2*p+2,j,IIP[j,2*p+2]):
> od: od:
> #--- contraction with u: t=u, tu=uu (TABLE V)
>
> for j from 0 to M do
> tu[j]:=uu[0,j]:
> od:
> for p to M do
> for j from 0 to J do
> IIIP[j,2*p+2]:=simplify(II[j,2*p+2]):
> print(2*p+2,j,IIIP[j,2*p+2]):
> od: od:
> for j from 0 to M do
> unassign('tu[j]'):
> od:
> #-----end-----

```

Frenet basis

A few comments are needed to explain how the Frenet curvatures are handled with the program.

We deal with the Frenet basis recursively. Our main objects are the scalars U_{ij} , the coefficients D_{ij} of expanding Frenet basis, $\vec{N}^{(i)} \equiv \sum_{k=0}^i D_{ik} \partial_\tau^k \vec{u}$ and the coefficients B_{ij} of the inverse expansion $\partial_\tau^j \vec{u} \equiv \sum_{i=0}^j A_{j,i} \vec{N}^{(i)}$. All these quantities are dealt with as functions of the Frenet curvatures κ_i and their derivatives.

Now, the knowledge of these quantities for all $i, j \leq p$ allows one to calculate them for all $i, j \leq p+1$ in the following way:

- Calculate $B_{p+1,p+1} = -\kappa_{p+1}B_{p,p}$. To prove this formula, one suffices to note that $\partial_\tau \vec{N}^{(p)} = \frac{1}{B_{p,p}} \partial_\tau^{p+1} \vec{u} + \dots$, which has to be expanded into $\kappa_p \vec{N}^{(p-1)} - \kappa_{p+1} \vec{N}^{(p+1)}$. However, $\partial_\tau^{p+1} \vec{u}$ contributes to $\vec{N}^{(p+1)}$ only. Therefore, $\partial_\tau \vec{N}^{(p)} = \frac{B_{p+1,p+1}}{B_{p,p}} \vec{N}^{(p+1)} + \dots$ and $\kappa_{p+1} = -\frac{B_{p+1,p+1}}{B_{p,p}}$.

- Calculate $B_{i,p+1}$ for all $0 \leq i \leq p-1$ using

$$U_{i,p+1} = \partial_\tau U_{i,p} - U_{i+1,p}$$

and $U_{p,p+1} = \frac{1}{2} \partial_\tau U_{pp}$.

- Calculate $B_{i,p+1}$ for all $1 \leq i \leq p$ using $B_{i,p+1} = -\vec{N}^{(i)} \partial_\tau^{p+1} \vec{u} = -\sum_k D_{ik} U_{k,p+1}$, the latter expression can be symbolically written as $\left\{ -\sum_k \frac{\partial \vec{N}^{(i)}}{\partial (\partial_\tau^k \vec{u})} U_{k,p+1} \right\}$. Note that $B_{p+1,0} = U_{0,p+1}$.
- Calculate $\vec{N}^{(p+1)} = \frac{\partial_\tau^{p+1} \vec{u} - \sum_{j=0}^p B_{p+1,j} \vec{N}^{(j)}}{B_{p+1,p+1}}$.
- Calculate $U_{p+1,p+1} = (B_{p+1,0})^2 - \sum_{i=1}^{p+1} (B_{p+1,i})^2$.